

**27.**

**Ocean Special Area Management Plan**

**Science Research Agenda**

**by**

**Monique LaFrance and Michelle Carnevale**

**University of Rhode Island Coastal Resources Center/ Rhode Island Sea Grant,  
October 2012**

Table of Contents

Executive Summary ..... 3

1. Introduction ..... 4

2. Research Focused on Baseline Data Collection ..... 8

    2.1 Observation stations..... 11

        Importance..... 12

    2.2 Physical oceanography..... 15

        Importance..... 15

    2.3 Wind and climatology ..... 16

        Importance..... 17

    2.4 Benthic habitat..... 17

        Importance..... 20

    2.5 Ecology ..... 21

        Importance..... 22

    2.6 Avian..... 24

        Importance..... 26

    2.7 Marine mammals and sea turtles (MM&ST) ..... 26

        Importance..... 28

    2.8 Fish ..... 28

    2.9 Fisheries (Commercial and Recreational)..... 31

        Importance..... 31

    2.10 Cultural and Historic Resources..... 32

        Importance..... 35

    2.11 Recreation and tourism..... 36

    2.12 Baseline research needs prioritized by location..... 38

3. Research Focused on Impact Assessment ..... 40

    3.1 Measuring impacts of offshore renewable energy (ORE) development ..... 40

        3.1.1 Transition of baseline studies into monitoring ..... 42

        3.1.2 National Oceanographic Partnership Program (NOPP) ..... 42

        3.1.3 Fisheries..... 45

        3.1.4 Electromagnetic fields (EMF) ..... 46

        3.1.5 Recreation and tourism..... 48

        3.1.6 Prioritization of monitoring research needs ..... 49

    3.2 Measuring the impact of climate change..... 50

4. Research Focused on Ocean Engineering ..... 51

5. Information Framework ..... 54

6. Regional Research Initiatives ..... 56

7. Conclusion ..... 57

References ..... 59

Appendix: Monitoring Protocols Developed by URI through the National Oceanographic Partnership Program (NOPP)..... 65

***Executive Summary***

With the State's adoption of the Ocean Special Area Management Plan (SAMP) as the primary planning and management document for offshore development, Rhode Island has moved to the forefront of marine spatial planning (MSP) in the United States. The Ocean SAMP Science Research Agenda (SRA) identifies where additional research is needed and recommends approaches for achieving these needs. The SRA focuses on the natural resources and the human activities occurring within Rhode Island's offshore waters, particularly the Ocean SAMP ecosystem.

The objectives of the SRA are:

- Assist in obtaining and directing future fundraising opportunities;
- Continue to learn about the Ocean SAMP ecosystem;
- Better understand the potential effects of future development, including offshore renewable energy (ORE), and other human impacts;
- Identify areas that warrant protection from certain human activities and areas for conservation;
- Increase Rhode Island's understanding of the projected impacts of global climate change;
- Obtain data at adequate spatial and temporal scales;
- Guide scientifically sound management decisions and support adaptive management strategies to minimize impacts to the environment and human activities;
- Promote the efficient development of ORE facilities where feasible; and
- Identify areas for collaboration and multidisciplinary research projects.

In order to fulfill the above stated objectives, the SRA proposes tasks regarding:

- **Baseline data.** Baseline data will permit further characterization of existing ecosystem patterns and processes, including environmental conditions and biological

distribution, abundance, and habitat use patterns, and also act as a reference point against which to measure change.

- **Monitoring.** Monitoring will allow for the assessment of impacts due to ORE development and global climate change. Assessment involves the identification and evaluation of the range and magnitude of potential effects associated with development or climate change on natural resources and human activities.
- **Advancing ocean engineering of offshore renewable energy devices.** Establishing appropriate design, construction, and structural requirements is essential to ensure an ORE facility reaches its maximum potential and efficiency.
- **Developing an information framework.** An information framework is a necessary component to organize, visualize, analyze, and disseminate information in a manner that is useful to all users (researchers, regulators, developers, public).
- **Supporting regional MSP efforts.** It is recognized some topic areas can be more fully understood over a broad geographic range (e.g. marine mammals, birds, fish). Additionally, expanding to a region focus encourages potential collaborations and funders for research initiatives.

The SRA was created with contributions from Ocean SAMP researchers, state and federal government agencies, environmental organizations, the public, and other stakeholders, including the fishing, recreation, and tourism industries. The development process was transparent, involving parties through questionnaires, as participants in meetings, and as document reviewers.

While the SRA focuses on the waters of the Ocean SAMP study area, the research ideas expressed may be applicable for regional MSP efforts or valuable on a national scale as it relates to understanding the effects of ORE development.

## ***1. Introduction***

The Ocean Special Area Management Plan (SAMP) process and resulting document has moved Rhode Island to the forefront of marine spatial planning (MSP) and offshore renewable energy (ORE) development at local and national scales. The success of the project was its

ability to accomplish extensive research goals and fully engage a variety of stakeholders throughout the two-year process. The Ocean SAMP greatly improved our knowledge of Rhode Island's offshore waters (Figure 1) through the compilation and synthesis of existing information and the collection of new data regarding the area's natural environment and human uses. The natural environment includes biological (fish, marine mammals, sea turtles, birds) and physical (seafloor and water column structures, wind field) components. Human uses refer development projects (ORE or other construction, dredging), recreation and tourism activities (boating, fishing, wildlife viewing), commercial fishing, and cultural and historical sites.

Since the completion of the Ocean SAMP, Rhode Island has continued to lead the way, further working to overcome the scientific and management challenges related to MSP and ORE development. The Science Research Agenda (SRA) presents an additional opportunity to address these challenges. The SRA has been created to identify and outline approaches for topic areas (related to natural resources, human uses, and engineering) that require further research within the Ocean SAMP area and regionally. Proposed research focuses on establishing baseline conditions, evaluating the individual and cumulative impacts of ORE development through monitoring studies, and addressing engineering challenges of ORE development. The SRA also works towards developing a standardized framework to organize and communicate research between and among users, including scientists, managers, developers, stakeholders, and the public. Although the SRA focuses on Rhode Island waters, the recommendations expressed may be applicable for MSP and ORE development efforts at other locations.

As an agenda, the SRA aims to satisfy the following objectives:

- Assist in obtaining and directing future fundraising opportunities;
- Continue to learn about the Ocean SAMP ecosystem;
- Better understand the potential effects, individual and cumulative, of future human activities, including ORE and other development, on the Ocean SAMP ecosystem;
- Identify areas that warrant protection from certain human activities;
- Better understand the potential impacts of global climate change;
- Obtain higher resolution spatial and temporal data;


- Guide scientifically sound management decisions and support adaptive management strategies to minimize impacts to the environment and human activities;
- Promote the efficient development of ORE facilities where feasible; and
- Identify areas for collaboration and multidisciplinary research projects.

Similar to the Ocean SAMP, studies proposed in the SRA were identified through contributions from a range of interested parties, including scientists, the Rhode Island Habitat Advisory Board and Fishermen’s Advisory Board, environmental organizations, stakeholders, and the public.

The need for additional research within the Ocean SAMP area was first recognized in the Ocean SAMP document (Section 1130. Applying Adaptive Management to Implement the Ocean SAMP):

“CRMC will develop and implement the Ocean SAMP Science Research Agenda, in coordination with the Ocean SAMP researchers, federal, state, and local government and other parties, to improve management policies and practices. The Ocean SAMP Science Research Agenda will allow CRMC to: 1) Continue to learn about Rhode Island’s offshore natural resources and human activities; 2) Better understand the potential effects of future development and other human impacts; and 3) Increase Rhode Island’s understanding of the projected impacts of global climate change. To develop the Science Research Agenda, the Council will put together an advisory group including scientists, partner federal and state agencies, environmental organizations, and users of the Ocean SAMP area. This group will help the Council to identify data gaps, short- and long-term research priorities, potential partners, and potential funding sources.”

The SRA is one mechanism to ensure the Ocean SAMP is implemented using an adaptive management approach. This management strategy has been undertaken by the Coastal Resources Management Council (CRMC) to manage Rhode Island’s coastal waters since its inception in 1971. Adaptive management is a systematic process for continually improving the management of an area by learning from the outcomes of previously employed policies and practices. This approach requires careful implementation, monitoring, evaluation of results, and adjustment of objectives and practices.



**Figure 1. Rhode Island Ocean Special Area Management Plan (Ocean SAMP) study area. The study area encompasses Rhode Island and Block Island Sounds and is approximately 1,500 sq mi.**

The SRA begins with a description of projects focused on providing a more complete baseline characterization for either the entire Ocean SAMP study area or particular portions of the study area (Section 2.0). The purpose of this research is to increase understanding of current physical and biological resources (conditions, patterns, and processes), as well as human use trends. To aid in directing research funding, section 2.0 prioritizes the research goals according to location: the entire Ocean SAMP study area, the Renewable Energy Zone (REZ), and the Area of Mutual Interest (AMI). Section 3.0 discusses projects aimed at monitoring and assessing the effects of ORE development, during construction, operation, and decommissioning phases. Section 3.0 also proposes studies needed to examine the potential effects of climate change. Section 4.0 discusses research related to advancing the engineering of ORE devices and facilities, dealing with design, construction, and structural concerns. Section 5.0 presents requirements for developing a framework to effectively organize, visualize, and distribute information. Lastly, Section 6.0 discusses how this Ocean SAMP SRA may relate to

other regional research initiatives such as the Northeast Regional Ocean Council (NROC) or the Northeast Sea Grant Consortium (NESGC).

## ***2. Research Focused on Baseline Data Collection***

The research proposed in sections 2.1 – 2.12 would greatly benefit our understanding of the Ocean SAMP ecosystem by providing the opportunity to further characterize baselines (environmental conditions and biological distribution, abundance, and habitat use patterns) and develop models. The studies should occur over multiple years to enhance temporal resolution of the data. To adequately consider human uses, baseline studies should occur throughout the Ocean SAMP study area within and outside of areas of interest for comparison purposes. Establishing outside “control” sites is necessary for detecting change, understanding causation, and providing evidence of impacts associated with human activities. While proposed for the Ocean SAMP area, the data needs described are presented for the consideration of any planning efforts, regardless of study location, to guide science and management actions.

Baseline studies involve the collection of data (physical, chemical, geological, and/or biological) to understand the current state of an ecosystem in its entirety or in part. To be most effective, baseline work is undertaken before a given project/use is initiated/permited. The current conditions act as a reference point, or baseline, which can then be used to compare subsequent data. Consequently, baseline data is critical for making well-informed, science-based management decisions, including those related to MSP.

For example, as a result of the baseline data collected during the Ocean SAMP process, conservation areas were identified; important habitats, foraging areas, and human use areas were recognized and documented as Areas Designated for Preservation (ADPs) and Areas of Particular Concern (APCs) (refer to Ocean SAMP, Chapter 11). Additional baseline data could be used to support the ecological importance of existing ADPs and APCs, or identify additional conservation areas/areas that should be protected from certain future ORE development or other disturbance activities. The Ocean SAMP policies direct potential developers to avoid APCs, and if avoidance is not possible developers must minimize and mitigate any impact to APCs. Current APCs include moraines, due to their high biodiversity, and some recreation and fishing

areas, due to their economic, ecological, and/or cultural importance. Alternatively, ADPs prohibit all development activities and currently comprise sea duck foraging habitat.

Baseline studies are more valuable when sustained over multiple years, as substantial inter-annual variability within the Ocean SAMP area has already been documented in nearly all processes and system attributes. Therefore, having detailed baseline data sets over long time series is essential to detect variability in ecosystem patterns and processes over various spatial (e.g. centimeters to kilometers) and temporal (e.g. intratidal, seasonal, annual, long-term) resolutions. Furthermore, these baselines are may be used to distinguish between changes due to natural causes, human disturbance (e.g. development, fishing), episodic events (e.g. storms, spills), and global climate change.

Baselines are often multidisciplinary in nature, which naturally supports ecosystem-based management (EBM) efforts. EBM considers the full array of interactions within an ecosystem between biology (benthic, pelagic), environment (surficial, sub-surface, water column), and humans (fisheries, recreation and tourism, cultural and historical resources). EMB and baseline studies foster collaborations between and among fields, encourage the collection and analysis of co-located and time-synced datasets, and permit the Ocean SAMP area to be examined on more comprehensive level and on an ecosystem scale. Collaborations can also improve the logistics of conducting research; for instance, vessel surveys could achieve various data collection goals to minimize expenses, time, and personnel.

As examples, primary production rates and phytoplankton blooms are often influenced by physical oceanographic parameters, such as temperature, stratification, and circulation; fish abundance and distribution patterns could be analyzed in conjunction with fishing activity patterns to provide a more complete picture of fisheries in the Ocean SAMP area; and underwater archeology could utilize much of the same data collected for benthic habitat characterization, physical oceanography, and fisheries studies. With respect to ORE, all disciplines can inform siting and assess impacts.

In addition, baseline data could be used to develop models and change scenario simulations of the Ocean SAMP area. Models and change simulations are powerful tools for:

- Producing realistic spatial and/or temporal portrayals of the ecosystem;

- Building an understanding of ecosystem attributes, patterns, and processes, and their relationships through the integration of data across disciplines;
- Identifying forcing mechanisms driving ecosystem processes, and then recreate situations and forecasts to confirm our understanding of such mechanism;
- Investigating hypothetical scenarios to forecast and further understand how changes (from human activity, natural variability, climate change) might alter the ecosystem or aspects of it;
- Identifying situations and areas for which permitting ORE development or other human activities is impractical;
- Suggesting hypotheses that can be tested with existing or new data; and
- Revealing data gaps.

To evaluate change, simulations with realistic forcing are compared to those with artificially modified forcing; models are executed under different scenarios by adjusting the input parameters to examine how the system would respond and the potential magnitude of the response. Consequently, modeling and change scenarios permit researchers to address specific questions about the ecosystem, such as those regarding ORE development and other human uses, global climate change, and natural variation. From a management perspective, these tools are valuable for guiding policy decisions and to prevent and/or mitigate issues.

As examples, models and simulations can offer more complex understanding of plankton blooms, primary production and respiration rates, temperature, salinity, wind strength and direction, and current and circulation pathways, the fates of waterborne materials (spill trajectories, dispersal patterns, and particle pathways), and the effect of turbine infrastructure on the seasonal stratification cycle. Moreover, physical oceanography model outputs could be valuable for numerous other studies (ecology, habitat, fisheries, and birds) and for deriving relevant ecological parameters (anoxia-hypoxia, stratification, tidally-driven patterns, current patterns). Currently, data collected for the Ocean SAMP report is being incorporated into a benthic habitat model.

**2.1 Observation stations**

*Establish a series of fixed buoy monitoring stations.* Observation stations would act as platforms for co-located *in-situ* acquisition of multidisciplinary data; the types of measurements that could be collected concurrently are shown in Table 1. Profile moorings are suggested, on which a single sensor package moves vertically and samples the entire water column. Profile moorings can capture vertical structure of the water column much more comprehensively, thus providing a better view of processes, including stratification and turbulent overturn characteristics. The data would be collected continuously to achieve high temporal resolution and high spatial resolution of the water column. Furthermore, profile moorings require only a single sensor package, making them more cost-effective than mounting several sensors at fixed depths.

**Table 1. Observation station data potentially acquired through fixed buoys.**

	<b>Data</b>
Biological data	Chlorophyll a concentration
	Primary production rate
	Respiration rate
	Fluorescence
	Nutrient concentrations
	Partial pressure CO <sup>2</sup>
	Total inorganic carbon (TOC)
	Phytoplankton and zooplankton abundance and distribution
	Food-web interactions
Physical data	Salinity
	Temperature
	Oxygen concentration
	pH
	Density

	Stratification
	Turbidity
	Turbulence
	Surface waves
	Current (speed and direction)
	Wind (speed and direction)
Marine Mammal data	Vocalizations

Observation station locations may include locations where development is proposed, such as the Renewable Energy Zone southeast of Block Island (REZ) and the Area of Mutual Interest (AMI) (Figures 2 and 3, respectively), to assess pre-development environmental condition. In addition, stations should be established at control sites where development is not set to occur for comparison purposes. Control stations should be located enough distance away from ORE sites to allow for the distinction of the effects of ORE development from natural ecosystem variability.

Successful examples of mooring-based efforts include programs in Long Island Sound, Narragansett Bay, and coastal waters (the Northeastern Regional Association of Coastal and Ocean Observing Systems [NERACOOS] and Mid-Atlantic Regional Association Coastal Ocean Observing System [MARACOOS]). These programs have proven influential in both the science and management arenas.

*Importance*

Establishing a series of fixed-buoy stations to collect co-located, multidisciplinary data would greatly benefit scientists and managers. These rich data sets would:

- Provide the opportunity for the Ocean SAMP area to be examined as an ecosystem as a whole, instead of fragmentary;

- Provide high temporal resolution data for baseline characterization of the existing ecosystem, recognizing and understanding change, and performing ecosystem-wide analyses;
- Improve spatial resolution of current data base
- Encourage investigations of relationships between and among different parameters. For example, physical oceanography measurements could assist in recognizing the causation behind primary production patterns witnessed in the Ocean SAMP study area. The great variability in primary production specifically makes the initiation of a longer term, multi-year data acquisition program essential in understanding ecosystem patterns and processes;
- Assist in assessing effects of development on the environment within and outside of infrastructure;
- Complement vessel-based surveys. Observation stations offer data at high temporal resolution, whereas vessel-based surveys offer high spatial resolution;
- Improve understanding of the physical processes driving the Ocean SAMP area and region; and
- Assist in developing and validating any models of oceanographic and climatologic parameters throughout the Ocean SAMP area. The combination of observation station and vessel-based survey data opens the potential for modeling parameters over suitable spatial and temporal scales.




Figure 2. Block Island Renewable Energy Zone (REZ).




Figure 3. Rhode Island and Massachusetts Area of Mutual Interest (AMI).

## **2.2 Physical oceanography**

Measure physical oceanographic parameters. Parameters to be measured are indicated in Table 1. The methods proposed to achieve this research goal are observation stations and vessel-based surveys. The establishment of a series of fixed buoy monitoring stations would provide high temporal resolution data and is described in section 2.1. Vessel-based surveys performed throughout the Ocean SAMP area would complement observation stations by offering data of higher spatial resolution. Surveys would collect vertical profiles of water properties using a CTD and would be conducted once monthly over a grid of 50-100 locations spanning the Ocean SAMP area. In addition, data could be collected off of the Block Island ferry; the ferry would be equipped with an ADCP to collect long duration repeat transect time series of vertical profiles of currents. The ferry could also carry nutrient and water quality sensors. Examples of similar vessel-based survey monitoring efforts, and their invaluable contributions to science and management programs, can be seen in Long Island Sound (Connecticut Department of Environmental Protection) and Narragansett Bay (Brown University; Atlantic Ecology Division of the US Environmental Protection Agency); these programs form a guide, for example regarding the kind of station spacing and survey frequency that would be most effective.

Develop physical oceanography models. Create a hydrodynamic/biogeochemical models to produce realistic simulations of the Ocean SAMP system. The user could then configure the input parameters (e.g. wind, temperature, salinity, transport rates, currents) and execute the model under different change scenarios to evaluate the system's reaction.

### *Importance*

The proposed physical oceanography data collection program is essential to:

- Characterize spatio-temporal patterns within the Ocean SAMP domain, with emphasis on full water column structure;
- Establish baseline conditions. It is recognized there is considerable variability of physical processes, such as current and circulation patterns, within the Ocean SAMP area, including inter-annually. Therefore, a long-term data collection program is needed to discern inter-annual trends and anomalies from climate change impacts;

- Recognize driving factors (e.g. wind forcing, estuarine exchange, shelf interactions, and tidal processes) behind observed variability and trends;
- Define circulation and transport pathways, including exchanges with nearby estuaries (Long Island Sound, Narragansett Bay) and the outer shelf, and exchanges between key constrictions (Point Judith and Block Island, Block Island and Montauk Point);
- Determine volume and rates of water transport;
- Identify budgets for water and abiotic/biotic water-borne materials; and
- Develop models and change scenarios.

### **2.3 Wind and climatology**

The unique combination of Rhode Island's water climatic and oceanographic conditions requires coupled monitoring of the wind field and sea-state. Such data would allow an accurate estimate the wind resource, and consequently optimize wind farm development within the Ocean SAMP area in terms of geographic location and footprint scale. To accomplish this goal, the following tasks should be undertaken:

Quantify the wind resource. The wind resource, outside the area of influence of the wind farm, can be represented by the mean wind speed distribution. An accurate estimation of the mean wind field can provide an accurate estimation of the expected power output from a single turbine at a specific location. However, the current spatial uncertainty in the wind field, in particular in the proximity of islands (such as Block Island), as well as vertical profile uncertainties, lead to potential expected power uncertainty, maybe as large as 30 % at hub height. Spatial and vertical monitoring of the mean wind field must be refined at the local scale using state of the art monitoring techniques, such as a floating (upward looking) LIDAR able to scan the vertical structure of the lower atmospheric boundary layer at a relevant vertical (up to 200 m in elevation) and spatial scale (order 200 m horizontal grid to validate AWS true wind model data).

Quantify complex airflow fields induced by a wind farm and the feedback on the wind field. Accurately estimating the extracted power from an offshore wind farm requires an assessment of the wind field between turbines. Since each turbine interacts with the incident wind field, the

wind field between turbines is modified and becomes far more complex than the undisturbed wind field outside of the wind farm. Accurately assessing this “inter-turbine” wind field requires complex modeling of interactions between the wind resource, turbines, boundary layer, and sea surface interactions (Moriarty, 2012). Such a model should be applied to the Ocean SAMP area, and in particular to the REZ, in order to assess the truly extractable power in Rhode Island’s offshore environment.

*Importance*

The proposed wind and climatology research is valuable to:

- Accurately define and predict the theoretical wind resource provided by the environment;
- Accurately define and predict the extractable wind resource in a wind farm; and
- Examine spatial and temporal variability of wind resource.


**2.4 Benthic habitat**

Mapping of priority areas. The Ocean SAMP focused on high-resolution mapping of areas identified as priorities for potential offshore wind development. This effort should continue in areas of high priority, as mapping provides valuable visual information of the geospatial distribution patterns and processes of numerous marine resources. The mapping process involves the collection (via vessel-based surveys) and integration of high-resolution geophysical data (side-scan sonar, bathymetry, and sub-bottom seismic reflection profiles) and ground-truth data (sediment and biology samples, underwater video and photography, sediment profile imagery). The types of maps that could be created are presented in Table 2.

**Table 2. Potential maps to be developed from future mapping projects.**

<b>Map</b>	<b>Purpose</b>
Quaternary geologic depositional environment	Present distribution patterns and extent of subsurface sediment characteristics, including the existence and thickness of stratified deposits (i.e. glacial lakefloor) and Holocene (marine) deposits. Identify extent and distribution of moraines (recognized as Areas of Particular Concern in the Ocean SAMP process)
	Help understand sediment transport and deposition processes

Biological communities/ species	Present distribution patterns and abundance of biological communities or species of interest
Benthic habitat	Examine and classify biological-environmental relationships through integration of data (e.g. benthic biology, water depth, depositional environments). Accomplished through statistical analyses. Good relationships between benthic geologic and biologic characteristics have been found previously within the Ocean SAMP area (LaFrance et al., 2010) (Figure 4)
	Present distribution patterns and extent of biologic-geologic characteristics of seafloor environments (Auster et al., 2009)
	Identify geological feature, biological communities/species of interest
Mobile fishing gear use	Identify locations utilized by finfish and shellfish trawlers from trawl scars visible in side-scan sonar records
	Provide use-intensity information
Cultural and historical resources (e.g. shipwrecks)	Survey data could lead to the discovery of unknown resources or confirm the location of known resources
	Aid in documenting fishing activity to identify important commercial and recreational fishing grounds



**Classification**

- 1.) *A. vadorum* (Type I) - Depositional Basin, silty sand
- 2.) *A. vadorum* (Type II) - Glacial Delta Plain, pebble gravel coarse sand
- 3.) *A. vadorum* (Type III) - Glacial Delta Plain, sheet sand
- 4.) *B. serrata* (Type I) - Glacial Alluvial Fan, boulder gravel concentration
- 5.) *B. serrata* (Type II) - Glacial Alluvial Fan, pebble gravel coarse sand
- 6.) *B. serrata* (Type III) - Glacial Alluvial Fan, sheet sand
- 7.) *J. falcata* - Moraine Shelf, boulder gravel concentration
- 8.) *Corophium* spp. - Moraine Shelf, pebble gravel coarse sand
- 9.) *P. remota* - Moraine Shelf, coarse sand with small dunes/sand waves
- 10.) *P. medusa* / *L. hebes* (Type I) - Glacial Alluvial Fan, coarse sand with small dunes
- 11.) *P. medusa* / *L. hebes* (Type II) - Inner Shelf Moraine, coarse sand sheets/waves/small dunes
- 12.) *Syllis* spp. / *P. medusa* - Glacial Alluvial Fan, sand waves
- Undefined

**Figure 4. Classification map of habitats offshore of Block Island. (From LaFrance et al., 2010, Ocean SAMP Technical Report 4). Each map unit, as defined by Quaternary geologic depositional environment type, is classified according to the most abundant genus. Environmental-biologic relationships were determined statistically.**

Validate and refine habitat template and ecosystem services typology models. It is not feasible (due to cost and time) to map the entire Ocean SAMP area at the resolutions required for micro-siting analyses and pre-permitting assessments; therefore, for broad scale characterizations, models must be developed and validated to provide information of the benthic ecosystem. The habitat template method (Kostylev and Hannah, 2007) is a model that uses measures of natural disturbance and physiological stress to estimate species diversity and community composition. These measures are modeled using broad-scale Ocean SAMP-area-wide datasets, including bathymetry, chlorophyll, and frictional velocity. The model would provide full coverage maps of the Ocean SAMP area and would be effective tools for understanding the dynamic of the ecosystem. In particular, they may be used to inform micro-siting and monitoring plans, and to model changes in diversity resulting from shifts in any of the model variables.

Once completed, the habitat template model must undergo extensive testing to confirm their predictions. Validation could be achieved by testing model results against actual data obtained from the Ocean SAMP process (including benthic macrofauna and fish sampling, underwater video, acoustic imagery; LaFrance et al., 2010, Malek et al., 2010), and from the collection of new data.

Develop indicators of seafloor disturbance. Indicators of seafloor disturbance could be species-based, community-based, or based on abiotic variables. Indicator development involves establishing baselines and expected biological responses, as has been done in Europe (Andrade et al., 2011, Rice et al. 2012) using current benthic sampling techniques. Indicators would be useful in the future for rapidly assessing the magnitude of impact at any stage of any project and for comparing the impacts of a wide range of coastal disturbances (oil and gas exploration, fishing, renewable energy projects) in the SAMP area and regionally.

#### *Importance*

- The proposed mapping research could direct future studies and guide management decisions, as it has the ability to:
- Provide baseline classification of habitats within the Ocean SAMP area against which to measure change over various temporal and spatial scales;

- Identifying features (geologic, fishing trawl scars, shipwrecks), critical use areas (essential fish habitat, foraging and nursery grounds, valuable fishing grounds), and areas of biologic importance (sensitive, high biodiversity, target biological communities or species) in need of further examination or that warrant protection from human disturbance;
- Identify Areas of Particular Concern (APC), such as moraines and other features with coarse-grained (gravel or larger) sediment;
- Use data, including APCs to direct construction locations by indicating where installation would be easiest and least expensive. For example, on Stellwagen Bank, geologic maps were used to plan a fiber-optic cable route, avoiding gravelly geologic habitats where the cable could not be buried for its protection. In addition, knowing the sub-surface depth to unstratified glacial deposits (till) and the semi-consolidated Cretaceous-Tertiary age strata could provide information about how turbines would need to be installed (pile-driving, drilling);
- Provide powerful modeling tools to aid in micrositing, guide monitoring, examine potential impacts of changes to the ecosystem due to environmental and anthropogenic factors, and identify ecosystem relevant spatial scales of sensitivity to those disturbances or changes; and
- Rapidly assess and compare the magnitude of impact of human disturbances using indicators of seafloor disturbance.

## **2.5 Ecology**

Measure ecological parameters. Data of interest are listed in Table 1 and includes primary production and respiration rates, nutrient concentrations, and phytoplankton and zooplankton dynamics. Data would be collected at observation stations and during vessel-based surveys using instruments such as plankton samplers, nutrient sensors, and optical sensors. The establishment of a series of fixed buoy monitoring stations would provide high temporal resolution data and is described in section 2.1. Vessel-based surveys conducted throughout the Ocean SAMP area would complement observation stations by offering data of higher spatial resolution. It is recommended these surveys occur once per month.

Develop models. Primary production and chlorophyll *a* measurements could be used to ground truth modeled results within the Ocean SAMP area derived from satellite imagery. Once the relationship between the two methods (*in situ* measurements and satellite data) has been established, satellite imagery could be used to increase spatial resolution of primary production measurements.

Measure benthic community organic matter consumption and nutrient regeneration. These measurements for the Ocean SAMP project were confined to areas of fine-grained sediments. However, sandy coarse-grained sediments dominate most of the Ocean SAMP area and organic matter consumption and nutrient regeneration cannot be measured using extracted cores (as done in the fine-grained areas). These sandy sites need to be studied using state of the art eddy correlation techniques.

#### *Importance*

The continuation of primary production measurements is vital in understanding how changes to the ecosystems may effect biological production. As the primary producers of coastal ecosystems, phytoplankton are responsible for supporting the growth of species at various trophic levels through the fixation of carbon and other required nutrients (Durbin et al. 2003; Pershing et al., 2005, Steele, 2001). Phytoplankton are the dominant prey item for zooplankton species and are integral in contributing to benthic-pelagic biogeochemical cycling and feeding benthic infauna as sinking organic matter and detritus (Durbin et al. 2003; Fulweiler et al. 2007; Smith et al. 2010). The timing and extent of the winter-spring phytoplankton bloom has been found to strongly correlate with fish larvae survival and recruitment success (Platt et al. 2003; Head et al. 2005). It is hypothesized that phytoplankton blooms contribute to the success of mature demersal fish by providing food for their prey items in the benthos, enhancing their condition, and permitting them to produce greater quantities and higher quality eggs (Friedland et al. 2008).


The proposed ecological research is necessary to:

- Perform a holistic examination of ecosystem function and biological production of the Ocean SAMP area;

- Produce baselines ecological parameters and their patterns, including primary production rates, organic matter consumption, nutrients, phytoplankton and zooplankton dynamics, and food-web interactions, over various spatial and temporal scales;
- Better understand primary production within the study area. Primary production rates can greatly fluctuate any given season or year depending on ambient conditions. The great variability in primary production specifically makes the initiation of a longer term, multi-year survey of primary production and chlorophyll a patterns in the Ocean SAMP area essential for creating a baseline of biological productivity for future management actions;
- Better understand driving forces behind and interactions between observed patterns;
- Calculate total system metabolism (production and respiration) and net air-sea exchange of CO<sub>2</sub> from partial pressure of CO<sub>2</sub> and total inorganic carbon. This would provide for the first time estimates of the excess production available to support higher level of the food web within the Ocean SAMP area;
- Better understand benthic community organic matter consumption and nutrient regeneration. The balance between the production of organic matter in the sunlit surface waters and the consumption of this organic matter in the water column and on the bottom provides the energetic base for fish and other higher trophic level production. The tight biogeochemical coupling of the water column and the bottom is a characteristic of virtually all coastal systems, though it is generally poorly document and quantified in areas with sandy sediments such as the Ocean SAMP area. The new techniques proposed, however, would provide insight of ecosystem characteristics; and
- Assist in developing and validating models of primary production rates from satellite imagery throughout the Ocean SAMP area.

## 2.6 Avian

Assess baseline distribution and abundance of avian species. Having additional baseline information about the spatial distribution and abundance of avian species would allow biologists a more detailed understanding of seasonal and annual variation in bird use within the Ocean SAMP study area. These data could also be used to identify key foraging and resting sites, migratory corridors, and important habitats used that should be protected from disturbance. This research should be conducted throughout the year to assess seasonal and annual variation in avian use of the study area. Surveys would be conducted along transects that are placed perpendicular to the avian density gradient (Figure 5; refer to Ocean SAMP Report for further details). Specific methodologies are discussed in Table 3.



**Figure 5. Avian aerial survey transects. (From Winiarski et al., 2011, Ocean SAMP Technical Report 11 addendum). Location of 24 aerial transects sampled from 2009 – 2012.**

Examine movement ecology of birds. Little is known about the movement ecology (flight altitudes, phenology of movements, movement direction) of birds in the Ocean SAMP study area, with the exception of satellite telemetry investigations of seabirds. This information

could be used to assess barrier effects and develop collision risk models for bird use in the study area. Methodologies to achieve this goal are discussed in Table 3.

Investigate factors affecting variability in avian distribution and abundance patterns. To better understand seasonal and annual variation in the spatial distribution and abundance of birds, information is needed on potential prey (fish, invertebrates, benthic organisms), habitat conditions (benthos, sea surface temperature, chlorophyll, salinity, fronts), and anthropogenic activities (e.g. recreational and commercial fishing, development). Having this data would lead to the development of models of distribution and habitat use patterns of avian species.

**Table 3. Methods for characterizing avian use.**

Method	Advantages	Execution
Ship-based line transect surveys	1.0 Provides spatially-explicit estimates of density 2.0 Allows identification of organisms to species-level, including threatened and endangered species 3.0 Access to areas close to Block Island	4.0 Transects similar to baseline studies using standardize protocols 5.0 Conducted throughout year to assess seasonal variation 6.0 Perpendicular to density gradient 7.0 Develop Density Surface Models for study area and adjacent buffer (Before-After- Gradient design) 8.0 Studies conducted before construction and up to 10 years after construction
Aerial –surveys (observer or videography)	<ul style="list-style-type: none"> <li>• Provide coverage over large areas over short time periods</li> <li>• Videography allows permanent record of detections</li> </ul>	9.0 Transects similar to baseline studies using standardize protocols 10.0 Conducted throughout year to assess seasonal variation 11.0 Transects located perpendicular to density gradient <ul style="list-style-type: none"> <li>• Develop Density Surface Models for study area and adjacent buffer (Before-After- Gradient design)</li> <li>• Studies conducted before construction and up to 10 years after construction</li> </ul>

Tagging (radio telemetry, satellite telemetry, GPS, nanotags)	<ul style="list-style-type: none"> <li>• Offers detailed data of movement ecology individuals</li> <li>• Records movement, and therefore, habitat use</li> </ul>	<ul style="list-style-type: none"> <li>• Tags would be tracked remotely (monitored by satellite, base stations, or directional antenna arrays)</li> <li>• Satellite telemetry would follow ongoing efforts with seabirds in RI</li> <li>• Studies prior to construction and up to 5 years post construction</li> </ul>
Land-based Radar	<ul style="list-style-type: none"> <li>• Provides flight altitudes, flight directions, and number of targets</li> <li>• Only protocol that can collect information on nocturnal migrants</li> <li>• Determine barrier effects and model collision risk</li> </ul>	<ul style="list-style-type: none"> <li>• Use a combination of vertical and horizontal X-band marine radar units</li> <li>• Operate 24 hrs per day 7 days per week, both pre- and post-construction</li> <li>• Use similar site on Block Island for pre- and post-construction surveys</li> </ul>

*Importance*

The proposed avian studies are important to:

- Characterize baseline avian distribution and abundance patterns
- Provide a basis to evaluate potential changes in avian use of the Ocean SAMP study;
- Data to could be used to assess barrier effects and develop collision risk models;
- Better understand current movement ecology by threatened and endangered species and develop models to assess potential impacts on sensitive species;
- Determine avian seasonal and annual use patterns;
- Identify key avian migration corridor, foraging grounds, habitat, and other important use areas; and
- Better understand factors affecting seasonal and annual variability in spatial distribution and abundance patterns.

**2.7 Marine mammals and sea turtles (MM&ST)**

Characterize spatial distribution, abundance, and seasonality of marine mammal and sea turtle species. The existing data assembled and summarized for the Ocean SAMP project provided relatively good background on the species diversity, abundance, distribution, and

seasonality of the marine mammals and sea turtles that occur in Rhode Island and Block Island Sounds. However, there is a lack of detailed, high-resolution information, in both spatial and temporal dimensions, for all species. For some species (smaller sea turtles, seals, and harbor porpoise), even the basic information is weak because those species are difficult to detect during surveys. This goal could be realized through one or a combination methods described in Table 4.

**Table 4. . Methods for characterizing marine mammal and sea turtle habitat use.**

<b>Method</b>	<b>Advantages</b>	<b>Execution</b>
Passive acoustic monitoring arrays	12.0 Provides continuous monitoring for the largest number of marine mammal species 13.0 Provides best broad-scale coverage 14.0 Can be used for any species that vocalizes	15.0 Arrays could be moorings, fixed bottom mounts, sensors attached to turbine bases, towed arrays 16.0 Sensors should be placed within and at decreasing distances from proposed development sites for comparison purposes 17.0 Increase sensors = increased coverage area and data richness
Vessel and/or aerial-based surveys	<ul style="list-style-type: none"> <li>• Provides coverage over large areas over short time periods</li> <li>• Can be used to collect new data or verify passive acoustic monitoring signals</li> </ul>	<ul style="list-style-type: none"> <li>• Surveys should be conducted at least biweekly</li> <li>• Surveys could follow currently ongoing surveys in an area of wind-energy interest south of Martha's Vineyard by a collaborative comprised of URI, New England Aquarium, and the Provincetown Center for Coastal Studies</li> </ul>
Tagging studies	<ul style="list-style-type: none"> <li>• Offers very detailed data of individual marine mammals or sea turtles</li> <li>• Records movement, and therefore, habitat use</li> <li>• Tags may also be equipped to record vocalizations of hosts</li> </ul>	<ul style="list-style-type: none"> <li>• Tags would be tracked remotely (monitored by satellite or by cellular telephone reporting)</li> <li>• Tagging protocols could follow those of NMFS for their programs in Massachusetts and Maine</li> <li>• Studies should begin at least one year prior to construction and continue throughout operation</li> </ul>

### *Importance*

Projects focused on marine mammals and sea turtles will be a necessary part of any marine spatial planning activities within U.S. jurisdiction because of the special protections afforded those species under federal legislation (Endangered Species Act, Marine Mammal Protection Act). The proposed marine mammal and sea turtle surveys are valuable to:

- Adequately understand marine mammal and sea turtle distribution and abundance patterns;
- Identify important use areas (seasonal, migratory pathways); and
- Provide a baseline to be used to detect and measure change.

It should be noted that the proposed research would not provide much data to address climate-change effects, since all of the marine mammals and sea turtles in the Ocean SAMP area represent only relatively small subsets of populations with ranges that encompass much larger regions—in some cases most of the North Atlantic Ocean.

## **2.8 Fish**

Identify seasonal habitat use by important fish and invertebrate species. A core challenge of developing spatial management plans is the acquisition of knowledge concerning the distributions, population structure, interactions and trends of key species and communities. To address this challenge, the RI Ocean SAMP sponsored a spatially explicit baseline survey of the demersal fish and invertebrate community in Rhode Island's nearshore waters (Malek et al. 2010). This survey was conducted primarily in the fall season to achieve synoptic sampling of the entire SAMP area. However, many fish and invertebrate species have seasonal migrations and therefore their distributions within the SAMP area differ throughout the year. A monthly otter trawl survey would capture distributional trends of the fish community throughout the year and provide a baseline against which to assess potential impacts of offshore developments. Furthermore, understanding spatial and seasonal patterns in the demersal fish community would allow for turbine construction to be scheduled when effects on vulnerable fish species are minimal. The continuation of a monthly otter trawl survey would provide a unique opportunity to monitor fisheries ecosystem change throughout all phases of wind farm development, from construction to operation to decommissioning. Research goals would be accomplished by

sampling a set of 6-8 fixed stations and 4-6 randomly selected stations within the Ocean SAMP area. Randomly selected stations would be stratified by depth and development interest to provide statistical power in future analyses. Sampling would be conducted monthly using a Rhode Island commercial fishing vessel equipped with a  $\frac{3}{4}$  scale NEAMAP-style net and a one inch knotless cod-end liner. Aggregate weights, counts, and individual length measurements would be recorded from all species collected, including lobster and other macroscopic invertebrates. Weather and sea conditions (i.e. wind speed, wind direction, and tidal stage) as well as water column profiles (water temperature, dissolved oxygen, and salinity) would be recorded prior to each tow to investigate the relationship between oceanographic conditions and the demersal fish community.

Sample hard-bottom habitats. The difficulty of towing gear on rough bottoms has resulted in limited sampling of hard-bottom habitats within the Ocean SAMP area and throughout the northeast region of the United States. Several important fish species (scup, sea bass, and tautog) and lobster are associated with hard-bottom environments, but their distribution and abundance are difficult to assess, as they are inadequately sampled. The proposed sampling approaches would be seasonal beam-trawl surveys, ventless traps, and fish pot surveys based on techniques established by the International Council for the Exploration of the Sea (ICES) Study Group on the Development of Fish Pots for Commercial Fisheries and Survey Purposes (SGPOT). The sampling strategy would consider factors like water depth, habitat type, and oceanographic conditions. In addition, a dedicated 7-10 day ROV cruise of moraines and other hard-bottom areas should be performed to further define these complex, high diversity habitats and to complement fisheries data.

Sample pelagic fish species. Pelagic fish are not well sampled despite that many species (herring, shad, menhaden, bluefish, striped bass) are both ecologically and economically important. To understand the distribution and movements of these species, sampling would be done through tagging and/or acoustic studies (Jolly and Hampton, 1990).

Identify fish food-web dynamics. During bottom trawl surveys, stomach samples and tissue samples would be taken from all target species and analyzed for diet composition and nitrogen stable-isotope signatures. These data would be used to discern food-web dynamics, including

predator-prey relationships and feeding history (Garrison & Link 2000; Hyslop 1980; Fry 1988; Hansson et al. 1997).

*Determine fish-habitat relationships.* Efforts to discern relationships between fish habitat and fish geospatial distribution and abundance patterns should continue. This research involves the mapping and sampling of benthic habitats and conducting fish trawl surveys. These data are then examined for statistically significant correlations to identify meaningful relationships of fish-habitat use patterns. Target species or entire fish assemblages may be evaluated.

*Importance*

The proposed standardized, long-term fish surveys and sampling programs would be useful to:

- Gain a more comprehensive understanding fish spatial and temporal distribution, abundance, and habitat use patterns within the Ocean SAMP area;
- Provide a baseline against which to detect and evaluate impacts, including from offshore development and climate change;
- Identify diverse and sensitive fish habitats that warrant protection;
- Establish seasonal fish abundance and habitat use to ensure appropriate timing of construction activities during the year;
- Investigate biogeographic patterns in predator-prey relationships through diet analysis;
- Quantify the trophic level of predatory species and further define time-integrated feeding history of each consumer using Nitrogen stable isotope analyses. This combination of techniques provides a spatially explicit model of species interactions;
- Establish multidisciplinary links, such as those between fishing patterns, fish distribution/abundance, and benthic habitats, and between seabird foraging, forage fish, and benthic habitats; and
- Establish protocols for effectively sampling hard-bottom habitats. There is a lack of fisheries studies sampling hard bottom, particularly in the northeast region of the

United States. There are numerous factors that go into sampling hard bottom, including pot design, bait type, string size, pot spacing, soak time, species specific catchability. This data gap presents a unique opportunity to work with fishermen in Point Judith to develop and test gear and methods for this type of work.

## **2.9 Fisheries (Commercial and Recreational)**

*Characterize fishing activity* Commercial and recreational fishing are some of the most economically and culturally significant human activities taking place within the Ocean SAMP area. Yet, there lacks adequate understanding of these activities with regard to when and where fishing occurs, species targeted, types of gear used, and economic data (e.g. commercial landings, fishing gear expenses, charter boat revenues). The Ocean SAMP fisheries chapter utilized existing data, including Vessel Trip Report (VTR) data and publically available landings data; while these data showed that commercial and recreational fishing are important within Rhode Island and Block Island Sounds and to the state of Rhode Island, the data presented are insufficient to understand changes to fishing activity as a result of factors such as ORE development and climate change. Therefore, additional information is required to improve understanding of fishing activity. Some of this information may be provided through existing data, such as gaining access and the ability to present data from Vessel Monitoring Systems (VMS) and log books from lobster vessels. However, not all fishing vessels are required to carry VMS, so there is still a need to ensure adequate representation across gear and vessel types. New quantitative and qualitative data describing fishing activity should be collected as well to supplement existing data sources. Such new baseline information could be collected through surveys and mapping exercises with fishermen. Furthermore, observational vessel-based surveys could be conducted, which could take place in conjunction with other research surveys (e.g. avian, marine mammal and sea turtle, fisheries, mapping).

### *Importance*

The proposed study would rectify the fisheries data gap by allowing researchers and management agencies to:

- Better understand the economic and cultural significance of commercial and recreational fishing within the Ocean SAMP area;

- Produce a more comprehensive understanding of fishing activity by examining spatial and temporal scales over which fishing occurs, as well as species targeted, types of gear used, and economic data;
- Establish a baseline of fishing activity to detect changes resulting from renewable energy or other projects, climate change, and/or changes to federal/state regulatory regimes. These results could then be tied to economic data on fisheries landings to evaluate the extent to which any economic effects felt by the fishermen could be contributed to different impacts;
- Identify areas heavily used by fishermen; and
- Establish links between fisheries, fish abundance and distribution patterns, and habitat to provide a more complete understanding of the Ocean SAMP area.

## **2.10 Cultural and Historic Resources**

Assess historic properties. The inventory and the identification of National Register-listed and -eligible historic properties is the critically important initial step in the Section 106 review process for federally funded, permitted, or reviewed undertakings, including ORE projects. Section 106 of the National Historic Preservation Act (“NHPA”) of 1966, as amended (36 CFR 800) requires federal agencies take into account the effects of their undertakings on cultural resources listed or eligible for listing in the NRHP (36 CFR 60). The agency must also afford the Advisory Council on Historic Preservation (“ACHP”) the opportunity to comment on the undertaking. The Section 106 process is coordinated at the state level by the State Historic Preservation Offices (“SHPOs”), which in RI operates within the offices of the RI Historic Preservation and Heritage Commission (“RIHPHC”). The issuance of federal and CRMC permits for offshore projects within the SAMP will depend, in part, on obtaining comments from the RIHPHC, as well as the comments of local tribal authorities (e.g., the federally-recognized Narragansett Indian Tribe’s Tribal Historic Preservation Office) and other interested parties (e.g., local historical societies and concerned citizens). The Ocean SAMP provided a broad contextual framework characterizing the cultural and historic resources within Rhode Island and Block Island Sounds, particularly for identifying potential shipwreck locations (Figure 6; Ocean SAMP Chapter 4). What is now needed is an assessment of historic properties.

This process involves a detailed examination and comprehensive inventory of cultural and National Register of Historic Places (“NRHP”)-listed and -eligible historic properties located both underwater and onshore in the coastal margin. Within the Ocean SAMP, cultural and historic resources primarily refer to shipwrecks and ancient Native American settlement sites and use areas. The proposed method to identify resources is vessel-based geophysical (side-scan sonar, multibeam echo-sounder, sub-bottom profiler) and imagery (underwater video and photography) surveys. The data is then reviewed visually for indications of resources. Once found, a resource needs to be identified and documented. In addition, determinations need to be made about excavation and protection. Currently, there is over 100 sq mi of geophysical data from the Ocean SAMP project that could be analyzed for archeological resources.





Figure 6. Potential historic shipwreck locations. (From Ocean SAMP Chapter 4).



**Figure 7. Sea level -50 m. Relative sea level was about 50 meters below present at 11,500 years before present (yBP). Brown colors represent dry land, blue represents water. Marine water is just impinging on the southern Ocean SAMP boundary. Other blue areas are possible lakes in closed depressions. The present-day shoreline is shown by the red lines. (From Ocean SAMP Chapter 4).**

Identify submerged paleolandscapes with archaeological sensitivity for containing ancient Native American settlements. Limited work has been performed to reconstruct paleolandscapes within Rhode Island and Block Island Sounds. At 11,500 years before present (yBP), relative sea level was 50 m lower, resulting in the shoreline being located at the southern Ocean SAMP boundary (Figure 7; from Ocean SAMP Chapter 4). It is this time period that the (currently) earliest evidence of human habitation of the northeast appears in the archeological record. From an anthropological perspective, understanding such paleolandscapes is valuable for making better predictions of where submerged landscapes may be, better understanding past human responses to changing environmental conditions, and helping to preserve the cultural history of Native Americans and their ancestors. From a scientific perspective, studies of past climatic and paleoenvironmental conditions may provide a more refined determination of sea-level rise. The proposed study involves geophysical surveying and collecting and analyzing core samples containing formally terrestrial strata.

*Importance*

The proposed cultural and historical resources research would be valuable to:

- Construct a comprehensive inventory of submerged and coastal historic properties (archeological sites, built resources, and traditional cultural properties) and cultural resources;
- Establish the necessary baseline data for assessing what impacts may result from development and other human activities;
- Better understand paleoenvironmental conditions within the Ocean SAMP dating back to the time of the last ice age;
- Better understand past climatic conditions within the Ocean SAMP, enabling researchers to discern long-term climatic trends from shorter term changes associated with human impacts;
- More precisely calculate the local rate of relative sea-level rise within the Ocean SAMP;

- Better understand past human responses to dramatic climatic changes and sea level rise through the identification and analysis of archaeological deposits of inundated ancient Native American settlements; and
- Increased knowledge about the archaeological resources within the Ocean SAMP may accelerate and simplify development project permitting processes.

As previous Ocean SAMP studies have indicated, cultural and historical resources “also contain ecological as well as cultural and historical information, and many are integrated into marine ecosystems as structures or as parts of the ocean floor environment” (Ocean SAMP Chapter 4). Underwater archaeological research in the marine environment is by its very nature multidisciplinary and collaborative, as much of the same data are collected for benthic habitat characterization (side-scan sonar, multibeam, sub-bottom profiler, underwater video, etc.), physical oceanography studies (water and sediment chemistry, temperature, current directions and velocities, etc.) and fisheries studies (correlations between fishing grounds, fishing methods, and the types, densities, and distributions of shipwrecks, impacts on shipwrecks from different types of fishing, etc.). These links are directly and indirectly applicable to submerged cultural resources identification, assessment and management.

### **2.11 Recreation and tourism**

*Characterize offshore recreational activities and tourism.* While it is acknowledged that the Ocean SAMP area is culturally significant for recreation and tourism (e.g. boating, sailing, wildlife viewing), there is limited understanding of the distribution of recreational activities. Questions, such as, “When and where are different types of activities occurring?” “How many boats are participating?” “What are size and types of these boats?” “Is the area being used primarily by Rhode Island residents or outside visitors?” need to be addressed to adequately understand the importance of recreation and tourism. Methods proposed to collect this data are public questionnaires and vessel-/aerial-based surveys conducted once monthly, which may occur in combination with other research surveys (e.g. avian, marine mammals and sea turtles, fisheries). Multi-year data would be necessary to account for annual variation (due to weather conditions, economic climate).

Quantify economic value of offshore recreational activities and tourism. The economic impact recreational and tourism activities in the Ocean SAMP area have on the state of Rhode Island is not well understood and the best available data are 10-20 years old (Tyrrell and Johnston 2001; Thunberg 2008). As examples, the only available data for Block Island Race Week dates back to 1986, and a state-wide study done by Ninigret Partners in 2006 found that the 43,000 boats registered in Rhode Island at that time generated approximately \$182 million worth of spending each year (R.I. Economic Monitoring Collaborative 2008). However, this figure excludes transients, megayachts (very large yachts), and regatta participants and therefore likely underestimates the economic impact of this industry. To improve our knowledge, the economic value of events (America's Cup, Block Island Race Week) and activities (recreational boating, sailing, wildlife viewing) need to be documented. The upcoming America's Cup events in the summer of 2012 presents such an opportunity.

Charter boat industry. The for-hire fishing industry is a significant recreational activity and economic driver within the state of Rhode Island. In 2009, there were 240 active party and charter boat licenses issued by the Rhode Island Department of Environmental Management. The waters off Block Island are heavily used by this industry because of the variety of different species found here throughout the year that are attractive to anglers. The proposed construction of five wind turbines off the coast of Block Island may have a variety of effects on both targeted fish species and on fishing activity, and may pose a threat to the viability of the party and charter boat industry. Before such development is permitted, a more complete understanding of the charter boat industry of the Ocean SAMP area, particularly around the Block Island REZ, is needed. Research should be undertaken to document and evaluate species catch levels, frequency of activities, spatial use, and economic value.

#### *Importance*

The proposed recreation and tourism studies are needed to:

- Identify temporal and spatial contexts at which recreation and tourism occurs, including for the charter boat industry;
- Identify types of activities that go on at what intensity;

- Provide a baseline of recreation and tourism activity, including for the charter boat industry, against which to measure change within the Ocean SAMP area; and
- Examine and quantify the potential impacts on recreation, tourism, and the charter boat industry from offshore development. Specifically, quantifying how activities contribute to the local economies of our coastal communities would provide regulators with a better understanding of the potential consequences of development and demonstrate the importance of considering these uses in management decisions.

The proposed research would benefit from a 5-10 year study period if offshore development occurs, since recreation and tourism activities would probably differ immediately following construction (interest may be generated towards viewing the turbines) versus over time (as turbines become established and excitement decreases).

### **2.12 Baseline research needs prioritized by location**

It is recognized that research priorities may vary based on location. The focus of some of the proposed research projects is to attain a more complete understanding the entire Ocean SAMP study area, whereas other research is specific to areas that may undergo ORE development. Table 5 summarizes what the research priorities are within three areas: (1) the entire Ocean SAMP study area; (2) the Renewable Energy Zone (BI); and (3) the Area of Mutual Interest (AMI). A description of the boundaries of each of these areas is provided in the Ocean SAMP document. These areas were chosen because they are likely to be the focus of any future funding opportunities. In some cases, areas represented as higher priority do not necessarily mean that no data is currently available; rather that additional data would be useful to better understand the resource.

From a management perspective, the AMI and BI are the highest priority areas to collect baseline data as they represent the two areas where future development may occur. Because the BI REZ has already been examined as part of the Ocean SAMP, baseline research needs for the BI study area are less extensive. Specifically, surveys of fish distribution and hard bottom habitat, identifying and assessing submerged cultural resources, and better characterizing charter and party boat activity represent the baseline studies that may be most informative for future management decisions. Alternatively, baseline data needs for AMI area are more

extensive due to the limited data collection conducted in the area during the Ocean SAMP process. The mapping of priority areas such as moraines, surveys of fish distribution and hard bottom habitat, identifying and assessing submerged cultural resources, and data on avian movement ecology and factors influencing distribution and abundance represent the highest priority areas for future baseline research. It is important to note that the research findings and lessons learned regarding effective methodologies gained through surveys of fish, benthic species and hard bottom habitats in the BI REZ may provide valuable insight into studies of the AMI area.

**Table 5. Baseline research needs prioritized by location.**

**“1” = first priority, “2” = second priority, and “C” = complete. \*Note the research goals identified in section 2 are primary overall needs, whereas this table is more spatially refined, prioritizing research goals based on study location.**

	Research needs for baseline characterization (pre- ORE development)	Study Area		
		BI	AMI	Ocean SAMP
Observation stations	Establish a series of fixed buoy monitoring stations for multidisciplinary data acquisition	1	1	2
Habitat	Mapping of priority areas	C	1	2
	Validate and refine habitat template and ecosystem services typology models	2	2	1
	Develop indicators of seafloor disturbance	2	2	1
Ecology	Measure ecological parameters	2	2	1
	Develop models	2	2	1
	Measure benthic community organic matter consumption and nutrient regeneration	2	2	1
Fish	Identify seasonal habitat use	1	1	1
	Sample hard-bottom habitats	1	1	1
	Sample pelagic fish species	2	2	1
	Identify fish food-web dynamics	2	2	1
	Determine fish-habitat relationships	2	2	1

MM&ST	Characterize spatial distribution, abundance, and seasonality	C	C	1
Avian	Assess baseline distribution and abundance	C	C	C
	Examine movement ecology of birds	2	1	1
	Investigate factors affecting variability in avian distribution and abundance patterns	2	1	1
Physical oceanography	Measure physical oceanographic parameters	2	2	1
	Develop models	2	2	1
Fisheries (commercial, recreational)	Characterize fishing activity	2	2	1
Cultural and historical resources	Assess historic properties	1	1	2
	Identify submerged paleolandscapes with archeological sensitivity for containing ancient Native American settlements	1	1	2
Recreation and tourism	Characterize recreational activities and tourism	2	2	2
	Quantify economic value of offshore recreational activities and tourism	2	2	2
	Characterize Charter boat industry	1	2	2

### **3. Research Focused on Impact Assessment**

#### **3.1 Measuring impacts of offshore renewable energy (ORE) development**

The recent interest in ORE within the United States has initiated concerns of impacts to the environment and on human activities. The development of suitable research programs and protocols to measure and monitor the effects of ORE is of national importance. Impact assessment of ORE involves the identification and evaluation of the range and magnitude of potential individual and cumulative effects associated with development (construction, operation, and decommissioning) on natural resources and human activities. Assessments are a vital step for any ORE project, providing evidence of impact occurrence, intensity, and sources. This information is critical for implementing appropriate adaptive management, mitigation, and conservation strategies.

ORE impacts are identified through monitoring activities. While baselines are important for understanding the current state and natural fluxes of an ecosystem, monitoring is useful for identifying deviations from that state. Similarly, just as long and detailed baselines are necessary to sufficiently understand biological and environmental patterns and processes, long-term monitoring is required to adequately characterize change and identify causal relationships. While change can occur due to human activities (ORE and other development, boating, fishing, etc.), global climate change, or natural variability and disturbance events, for the purpose of this section (3.1) of the SRA, changes refer to those as a result of ORE development.

Models and change scenario simulations derived from monitoring data can be valuable tools prior to any decision making and further support adaptive management strategies. Models/simulations can:

- Allow for different development scenarios to be weighted against one another (by comparing model/simulation results);
- Provide insight into how alterations to the environment may influence different aspects of the ecosystem and/or the ecosystem as a whole;
- Allow individual and cumulative impacts of ORE development to be assessed;
- Increase understanding of an area (attributes, patterns, and processes) over suitable spatial and temporal scales; and
- Promote investigations of data trends and relationships.

The sections below discuss two recommended monitoring strategies for assessing the effects of ORE development on the marine environment and human activities. The first strategy (3.1.1) is monitoring via the extension of the baseline studies described in section 2. The second strategy (3.1.2) focuses on methodologies to detect, measure, and monitor specific effects of ORE development. For both strategies, studies should be implemented throughout the Ocean SAMP region within and outside of areas proposed for ORE or other development. Having undeveloped “control” sites permits comparisons to be made with developed sites and facilitates better understanding of development impacts. With regard to duration, the proposed monitoring

research should be sustained over a period of years to capture variability and sufficiently understand initial, short-term, and long-term impacts to the area.

*3.1.1 Transition of baseline studies into monitoring*

Monitoring activities to assess impacts of ORE development may be structured as a continuation of baseline studies (discussed in Section 2). Rather than be used to establish baseline conditions, these studies would transition into a monitoring program. This would allow for 1.) Direct comparison of baseline and monitoring data over time; 2.) Changes (physical, biological) within the Ocean SAMP ecosystem to be more readily detected and fully understood, with a focus on those due to ORE development; and 3.) More accurate development of models and change scenarios.

*3.1.2 National Oceanographic Partnership Program (NOPP)*

Through its involvement in the National Oceanographic Partnership Program (NOPP), the University of Rhode Island (URI) is currently leading the monitoring-based project “Developing Environmental Protocols and Modeling Tools to Support Ocean Renewable Energy and Stewardship.” This project has further placed Rhode Island in the forefront in the United States for identifying and assessing development impacts of ORE facilities on marine resources and human activities, and reports generated with provide a national service. This NOPP project is two goals. First, the project aims to identify potential effects of ORE at various development scales (demonstration, commercial, multiple commercial), indicate the anticipated intensity of those effects, and suggest where additional data is needed. The second goal of the project is to develop comprehensive protocols for measuring and monitoring the effects of ORE development (Table 6). To accomplish these tasks, URI has completed an exhaustive literature review of nationally and internationally published work related to ORE, and is now in the process of assembling and evaluating methods to produce the most effective suite of monitoring protocols. Summaries of the protocols listed in Table 6 are presented in the Appendix.

**Table 6. NOPP protocols for measuring effects of ORE development. Cultural and historic resources are discussed in NOPP report 1.4. The remaining topic areas are discussed in NOPP report 1.5.**

<b>NOPP protocols for ORE impact assessment</b>
<b>Benthic Resources and Habitat</b>
1. Scour and/or deposition
2. Changes in Benthic Community Composition
3. Reef Effect (Increase in Hard Bottom Habitat)
4. Changes in Hydrodynamics
<b>Avian Species</b>
1. Ship-based Visual Surveys
2. Aerial Surveys using Human Observers
3. Aerial Surveys using High Definition Videography
4. Aerial Surveys using Digital Still Photography
5. Radar Surveys
6. Visual Surveys of Flight Ecology
7. Flight Call Surveys
8. Using Radio Telemetry to Assess Movements
9. Using Satellite Telemetry to Assess Movements
10. Using GPS Tracking to Assess Movements
11. Thermal Animal Detection System to Assess Collision Risk
12. Sonar and Video Technology
<b>Marine Mammals and Sea Turtles</b>
1. Visual Surveys
2. Passive Acoustic Monitoring
3. Marine Mammal Observers

4. Stranding Response Networks
5. Tagging
6. Stress Hormone Assessment
7. Underwater Photography
8. SCUBA Surveys
9. ROV Surveys
<b>Fisheries Resources and Fishing Activity</b>
1a. Trawl Surveys
1b. Ventless Trap Surveys
2. Monitoring for Project-Scale Changes
3. Reef and Aggregation Effects
4. Blade Strikes
5. Spatial Use of Fishing Activity
<b>Cultural and Historical Resources</b>
1. Geophysical Surveys
2. Cultural Landscape Approach (CLA)
3. Archaeological Sensitivity Analysis (ASA)

A wind facility consisting of 5-8 turbines is being proposed within the Renewable Energy Zone (REZ) off of Block Island. This proposed demonstration-scale project would present a unique opportunity to directly detect, assess, and monitor impacts on marine resources and human activity in the immediate and surrounding environment of the ORE facility. While it is recognized a demonstration-scale project may not reveal all effects of larger-scale ORE developments, it would provide a strong starting point.

The proposed Block Island ORE project would be valuable to:

- Better understand potential discrete and cumulative effects of ORE on natural resources and human activities within the Ocean SAMP area;
- Design monitoring protocols and establish programs for assessing specific and cumulative impacts of ORE development;
- Guide adaptive management strategies. As an impact is being monitored, policies can be enacted to mitigate issues and/or prevent future occurrences;
- Validate and refine models and change scenarios of the Ocean SAMP area developed from baseline data; and
- Promote collaboration with research groups and management agencies to design standardized monitoring protocols and cumulative impact assessment tools in conjunction with one another to increase their overall compatibility and effectiveness, and reduce research costs and time.

### *3.1.3 Fisheries*

The for-hire fishing industry is a significant recreational activity and economic driver within the state of Rhode Island. The construction of five wind turbines off the coast of Block Island may have a variety of effects on both targeted fish species and on fishing activity, and may pose a threat to the viability of the party and charter boat industry. During construction, party and charter vessels will be displaced from the vicinity of construction activity. Construction noise along with vessel activity may cause fish to leave the area, decreasing catchability. Furthermore, little data exists to suggest what changes to fish distribution and abundance may occur once turbines have been constructed and are operational. Some of the species targeted by the party and charter boat industry may move elsewhere in response to habitat disturbance, whereas other evidence suggests the placement of new structures in the marine environment will result in reef effects as benthic organisms growing on the turbine foundations attract various species seeking food or shelter. In the Gulf of Mexico, reef effects have created habitat for numerous recreationally-targeted fish species on the jacketed structures of oil rigs, and in return these oil rigs serve as important fishing grounds for recreational fisheries. The jacketed structures of the wind turbines planned for construction off Block Island may have similar reef effects. However,

fish species most commonly targeted by the party and charter boat fleet are pelagic species, including striped bass and bluefish, which may not be attracted to new structures in the water.

The above-mentioned potential effects to party and charter boats could also have secondary economic effects. For example, if during the construction period, vessels are forced to travel further, or to areas where fish are less abundant, this could have negative economic consequences for their businesses. Likewise, if the most desirable species become more difficult to catch on a short-term or long-term basis, a loss of clients paying for fishing trips could result. Conversely, if reef effects concentrate or attract fish, and increase catchability, wind turbines could eventually be seen as a boon to the industry.

In partnership with the Rhode Island party and charter boat industry, we propose to determine the changes in activity and catch level of the party and charter boat industry during pre-construction, construction, and post-construction periods within and surrounding the Ocean SAMP REZ (Figure 2). This data will allow researchers to: 1) develop a baseline for where fishing is taking place; 2) determine effects of construction on spatial use of the area by the party and charter boat industry; and 3) document activity changes within waters around Block Island post-construction.

We also propose to analyze economic effects of the potential changes to fishing activity as a result of the construction and operation of the Block Island wind farm by documenting changes in: 1) the number of paying clients during this time period; 2) the number of return clients; and 3) the changes to the cost of fishing during this time.

NOPP fisheries protocols 1, 2, and 5 (see Appendix) directly address these concerns.

#### *3.1.4 Electromagnetic fields (EMF)*

The effects of EMF emitted from submarine cables on marine organisms is largely unknown and of high concern to commercial and recreational fishermen in RI, as well as other fishermen throughout New England. The Block Island wind farm project presents the opportunity to directly address the dearth of knowledge on EMF and clarify how fisheries throughout New England may be impacted. The project focuses on determining the actual emissions from cables associated with offshore wind farms and any response by sensitive receptor organisms. The proposed cost is \$2.3 million over 5 years.

A two strand approach is proposed: 1. Field surveillance monitoring (ambient and cable related); 2. Experimental studies to assess the response of receptor organisms. To accomplish the first objective, the ambient EMF would be measured through deployment of a dedicated ROV with EMF sensor. The EMF-ROV will be able to measure existing sources of EMF (e.g. subsea structures, geomagnetic field), as well as the magnetic field component from energized cables. It is this magnetic emission from both DC and AC subsea cables that is within the range of detection by receptor organisms. In addition, the magnetic field induces electric fields in the seawater and seabed as a result of water movement and/or organism movement through the field (i.e. electromagnetic induction). We anticipate the sensor can be moved around the wind farm development and to control sites and either survey along transect lines (or cables) or be stopped at sample points over the area to collect data from a static position, thereby enabling EMF decay to be quantified (rather than relying on assumption-based modeling).

To complement the field based surveys and to provide the actual evidence of whether the EMF emitted are detectable by receptor organisms, and of any importance when considering environmental impact we suggest a program of field and laboratory experiments. In the field, baited cameras would attract commercially important fish and/or crustaceans. We would then quantify behavioral response next to cables and compare with control studies away from the cable, but otherwise in similar environment conditions. An important aspect of these behavioral responses will be ensuring that the sampling size will be statistically sufficient to determine emergent properties from multiple individuals to look for higher scale effects (i.e. population level impact). As a precursor to the main studies with the baited cameras we have planned laboratory experiments to quantify the type of response that each target species had to controlled EMF of different levels. The quantified behaviors would then be used to define the type and level of response seen in the baited video studies. The laboratory and field based studies have been planned according to appropriate experimental design and the measured variables would be focused on determining thresholds of response by the target species and clear quantification of species specific responses.

Lastly, a tagging study is proposed to better understand the effects of EMF on commercially important crustacean species, mainly the American Lobster (*Homarus americanus*). Because it is well understood that off Rhode Island the American Lobster naturally migrates from waters

offshore into Narragansett Bay, the movement of lobsters over installed cables may be tracked. The proposed mark and recapture study would catch and tag lobsters from Narragansett Bay working in coordination with local lobster fishermen, relocate them to areas across the cable route, and measure the recapture rate before and after cable installation. The results of this study will help inform fishermen and regulators on what the actual effects of EMF are on this important fishery.

### *3.1.5 Recreation and tourism*

The proposed study aims to measure changes in recreational boating activity and the perceptions of tourists to Block Island as a result of offshore wind turbines in the REZ off Block Island. Block Island is an important driver for tourism in the state of Rhode Island, attracting over half a million visitors per year (Global Insight, 2008), including recreational boaters. The construction of five wind turbines off Block Island's shore adds an unknown element to the island's tourism industry. The novelty of turbines may attract tourists or recreational boaters who wish to view them; on the other hand, the new turbines may negatively influence perceptions of tourists attracted to the island's relatively unspoiled views. Moreover, during construction recreational boaters will likely be excluded from the project site for safety reasons. The Ocean SAMP recognized this potential effect, particularly around Block Island; Section 1160.9, Monitoring Requirements #2 stated:

“The Council shall require where appropriate that project developers perform systematic observations of recreational boating intensity at the project area at least three times: pre-construction, during construction, and post-construction. Observations may be made while conducting other field work or aerial surveys and may include either visual surveys or analysis of aerial photography or video photography. The Council shall require where appropriate that observations capture both weekdays and weekends and reflect high-activity period including the July 4<sup>th</sup> holiday weekend and the week in June when Block Island Race Week takes place. The quantitative results of such observation, including raw boat counts and average number of vessels per day, will be provided to the Council.”

With regard to the perceptions of tourists visiting the island, qualitative surveys will be used. Tourists will be asked to evaluate what they view as the positive and negative impacts of the construction and operation of the Block Island wind farm, including what they believe are

potential environmental effects. Surveys will be conducted and responses analyzed before and during construction, and during operation of the facility to determine how perceptions change during the different phases of the project.

Changes in recreational boating will be quantified through weekly visual surveys recording boat counts and types of activity occurring at the project site and two other sites (controls) around the island during summer months (May – September). These surveys should be conducted at least one year before construction begins, during the construction of the facility, and for multiple years once the project is operational. Surveys will be conducted using a methodology developed at the University of Rhode Island by Thompson and Dalton (2010). Data will be analyzed to assess changes in the number of boats utilizing the project site and the control sites. Geospatial analysis will also be conducted to evaluate whether boating activity is being shifted to other areas around Block Island. Because the project may potentially span up to 5 years, this study will be able to measure both immediate changes caused by construction activities and longer-term impacts to recreational boaters at/near the project site. Changes in intensity and types recreational boating activities will also be evaluated and analyzed to determine the economic impact such of changes.

### *3.1.6 Prioritization of monitoring research needs*

Table 7 below identifies monitoring studies of highest priority for assessing impacts of ORE development (during construction, operation, and decommission phases). Primary impact concerns are fisheries (characterized by changes in fishing activity and catch levels) and marine organisms (signified by changes in species distribution and abundance). Also of high concern with regard to marine organisms are monitoring and evaluating the effects of noise associated with ORE development on marine mammals, EMF from cables on elasmobranchs and the American Lobster, and avian collision rates with ORE devices. High priority studies to assess the physical environment include generating noise profiles for ORE during all phases of development and establishing fixed observation buoys to collect physical oceanographic data. The priority research goals related to recreation and tourism include monitoring changes in recreational boating, tourism activity, and public perception.

Table 7.

	<b>Research needs to monitor and assess impacts of ORE development</b>
EMF	Effects on marine organisms (specifically the American Lobster and elasmobranchs)
Acoustic	Understanding of construction, operation, and decommission noise associated with ORE
	Impacts of ORE noise on marine organisms (specifically marine mammals)
Observation stations	Establish a series of fixed buoy monitoring stations for multidisciplinary data acquisition
Habitat	Changes in benthic community composition
Fish	Changes in species distribution, abundance
Avian	Collision rates of species
	Changes in species distribution, abundance
Fisheries (commercial, recreational)	Changes in fishing activity
	Changes in catch levels
Recreation and tourism	Changes in recreational boating and tourism activity

### 3.2 Measuring the impact of climate change

Marked climate change impacts have been/are being seen in shallower systems adjacent to the Ocean SAMP area (Narragansett Bay and Buzzards Bay) (refer to RI Ocean SAMP document, Chapter 3). Over the last century, air and sea temperatures, precipitation, sea level and cloudiness have increased in New England (Smith et al 2010; Nixon, 2009). Annual chlorophyll biomass has also decreased dramatically (from 1970s-2006) and winter-spring phytoplankton bloom biomass and has decreased or in some instances not occurred (Oviatt 2004; Oviatt et al. 2002). With the weakening and changing phenology phytoplankton blooms,

it is hypothesized that the rate of re-mineralization of nutrients in the benthos will concurrently change due to weaker benthic-pelagic coupling and that less organic matter deposition will result in fewer secondary consumers and ultimately less demersal fish (Fulweiler and Nixon 2009; Nixon et al. 2009). While research over the last 60 years have elucidated some of the correlations between the changing physical environment and phytoplankton production in Narragansett Bay, very little is known on how Block and Rhode Island Sounds have responded to the same climate trends, including with regard to primary production, respiration, and coupling between the seafloor and water column. Therefore, the time frame for the proposed baseline and monitoring research (refer to sections 2 and 3, respectively) should be extended. This continued data stream would build the knowledge platform necessary to document, detect, and interpret climate change impacts within the Ocean SAMP area over spatial scales ranging from ecosystem-wide to individual habitats.

#### ***4. Research Focused on Ocean Engineering***

Ocean engineering is a critical component of optimal offshore renewable energy (ORE) development. Advancing the engineering of ORE devices and facilities is essential for establishing appropriate design, construction, and structural requirements to:

- Ensure an energy facility reaches its maximum potential and efficiency;
- Adequately anticipate individual and cumulative effects of ORE development;
- Use understanding of ORE effects to identify areas to be protected;
- Minimize the time and costs of project construction, operation, and decommissioning;
- Address safety concerns;
- Recommending design and structure standards; and
- Inform the Certified Verification Agent (CVA) review process used by CRMC and the Bureau of Ocean Energy Management (BOEM).

Therefore, advancing the engineering of ORE is valuable not only for the Ocean SAMP area, but nationally and globally, as well.

URI has submitted a proposal to the US Department of Energy (DOE) in partnership with developer, Deepwater Wind, turbine manufacturer, Siemens Inc., and other academic institutions (National Renewable Energy Laboratory, Massachusetts Clean Energy Center, and the State University of New York). The proposal is to study the demonstration-scale project proposed within the Ocean SAMP (BI) REZ for designs and techniques to minimize uncertainties, time, and costs associated with energy production from offshore wind farms. Through monitoring and modeling of the five 6MW turbines and associated support structures (jacket structures, pile foundations, submarine transmission cables), the project aims to: 1.) Conduct extensive multi-disciplinary monitoring and modeling of the BI wind facility, to optimize the construction, deployment, and operations of such projects in the future, leading to a reduced cost of energy; and 2.) Use data acquired and lessons learned to reduce the deployment timelines and uncertainties for larger commercial scale offshore wind development projects in US waters.

The following is research that should accompany a demonstration-scale wind facility:

- *Observation buoys.* Design, install, and operate a buoy based observation system to make measurements of key physical parameters to characterize and understand environmental forcing of the turbine structure including waves, currents, winds, and salinity and temperature in the immediate vicinity of the turbine structures. Wind characterization will include measurements with elevation via buoy mounted LIDAR. This data would provide the first comprehensive data set and examination of the resulting structural and turbine response of a full-scale jacket support structure to extreme environmental forcing (wind, wave) and cyclic (fatigue) loads. Such information would be invaluable to the industry for optimizing the support structure and the layout designs of a wind farm, including to minimize wake effects.
- *Instrument the wind turbine foundation system.* Place instruments on the foundation system to measure cyclic stresses, accumulated displacement, and pore water pressure generation in the surrounding soils. Strain gages on the piles will monitor the stresses and accumulated displacement and in situ piezometers will measure the generation of pore pressures. In the glacial silts and sands in Rhode Island Sound,

possible generation of pore pressures can lead to a reduction of soil stiffness under long-term cyclic loading.

- *Turbine structure monitoring.* Implement a structural health monitoring system of the lattice jacket structure by placing strain gages and accelerometers on the jacket and analyzing the data periodically and after extreme events (i.e. hurricanes) to predict on-going and future structural damage. This would help with the configuration of an optimal design for fatigue based on accurate measurements of stiffness and damping of the structure/foundation system.
- *Models of turbine response to environmental loading.* Develop and implement models to predict the environmental loading and response of the structure and foundation to routine and extreme loading. Compare the model predictions to data collected from the observation buoys, instruments placed on turbine foundation system, and monitoring.
- *Passive acoustic monitoring system.* Design and implement a passive acoustic monitoring system to monitor ambient noise levels. The system would measure ambient noise levels prior to construction, as well as noise generated during construction, operation, and decommissioning. The measurements should allow determination of noise associated with marine biologics, marine shipping, and turbine construction (pile driving) and operation. Adding a modeling component will develop techniques for predicting the propagation of radiated noise from the structure during construction (pile driving) and operation. Model results will be compared and validated with acoustic measurements during construction and operation.
- *Assessment of energy production.* Perform an analysis of the power production and energy transfer/integration into the grid of the selected turbine by wind direction, atmospheric stability, and wind speed and shear. Determine the capacity factor for the turbine, assess possible wake effects from adjacent turbines in the field, and compare estimates of the predicted to observed power production.
- *Costs of energy production.* Modeling investigations and observations would allow for comprehensive offshore wind economic modeling and analysis to compute costs

of energy production under various scenarios. This effort would take into many factors into consideration, including a better understanding of the efficiency of turbines in a wind farm, wake effects, and true structural/foundation response due to environmental loads.

- *Improve design standards.* Based on modeling investigations and observations, perform an in depth review of the current design standards of the wind turbine support structure and foundation. Then, recommend how standards should be revised to ensure high performance and efficiency of turbines and more accurately represent the risks of ORE development. Establishing engineering requirements for ORE projects within the United States is an area of active interest, for which the proposed research could offer valuable information.
- *Certification Verification Agent (CVA) review process.* Advancing the engineering of ORE and having a greater understanding of ORE design standards would improve the CVA review process required by locally and nationally by CRMC and BOEM, respectively. The CVA process is described in chapter 11, section 1160.6 of Ocean SAMP document.

## **5. Information Framework**

Adequate geospatial support for the Ocean SAMP area involves the development of a framework to facilitate organization, visualization and dissemination of information, as well as guide policy decisions and monitoring. Geospatial data and their relationships are the cornerstone of marine spatial planning (MSP); whether preparing for offshore renewable energy (ORE) or assessing vulnerability to climate change. Geographic Information Systems (GIS) provide the suite of tools to analyze and display these relationships in a manner that is useful to both researchers and regulators. Integrating resource use with scientific studies detailing the dominant biological, physical, social and ecological relationships provides managers the knowledge to develop strategies that provide needed access to resources without compromising environmental conditions. Tasks for this work are outlined below.

*Data integration and consolidation.* The central database developed as part of the Ocean SAMP project is the core structure for integrating and visualizing all of the research conducted

throughout the 2-year process, and it is imperative that any new information collected is compatible with the existing system. The Ocean SAMP data holdings are the only multidisciplinary data source available for Rhode Island's offshore waters.

*Mapping and data exchange.* Access to Ocean SAMP data currently is done through the NARRBAY.ORG website where users can download individual data sets or access data through the RI Ocean SAMP Map Viewer. These delivery outlets allow for the efficient mapping and data exchange between and among scientists, managers, and the public.

The NARRBAY.ORG website was modeled after the RIGIS website, the primary point of access to terrestrial geospatial data for Rhode Island, to access Rhode Island's coastal and marine-related data, and was developed to accommodate the rich variety of file formats. The software architecture of the NARRBAY.ORG site is now 10 years old and in need of a technology makeover. Newer technologies, such as content management systems, are now available that would make the website more efficient and easier to maintain. A redesigned NARRBAY.ORG site would be fully searchable and would integrate directly with the RIGIS system for cross-platform querying. Along with providing direct data download, the site would feature the latest news feeds for marine spatial planning and ORE development, and would house links to related regional organizations and data providers.

In addition, the Environmental Data Center (EDC) at URI would leverage new efficiencies in cloud computing found within the ArcGIS.com platform, rather than develop a single map viewer, for visualizing Ocean SAMP data. The system is designed specifically for spatial data and is a cloud-based, content management system for working with geographic information. The platform provides an on-demand infrastructure for creating web maps and collaboratively sharing that information throughout an organization. There are many benefits of this model. For instance, users would be able to search the available data and build their own custom maps using the standard Ocean SAMP template, which could be saved and shared to improve collaboration. As new data are acquired they would simply be added to ArcGIS.com for immediate access.

*Information access and integration within management agencies.* Spatial relationships are fundamental to resource management; who, what, when, where and how the coastal

environment is being used are all issues that factor into management decision-making. A GIS provides the suite of tools to analyze and display these relationships, but there is a cost in terms of the time required to obtain/maintain proficiency with the software. Oftentimes individuals accumulate some level of experience on project-specific basis, but with limited funds there is often not enough time available for a thorough review of how information is stored and disseminated throughout an organization. To mitigate these issues, the EDC is proposing to expand the in-house mapping capabilities of the CRMC, work with staff to document the GIS workflow and needs for different focus areas, and provide technical training. The implications of doing so extend far beyond the Ocean SAMP project and would continue to benefit the organization and its mission long after the immediate project has ended.

*Regional collaboration.* The EDC will engage appropriate regional organizations to share maps, data, and information, and encourage regional collaboration on MSP of shared coastal resources. The regional collaboration effort will target groups with missions that embrace open sharing of data and experiences.

## **6. *Regional Research Initiatives***

Regional research initiatives in many ways expand upon and compliment the research projects presented here or the research already conducted as part of the Ocean SAMP process. Regional initiatives will result in a refined understanding of both the Ocean SAMP and New England areas by providing information over a broader geographic range. Many ecosystem attributes and human uses, such as the distribution of marine mammals, pathways of migratory birds, and fisheries activities, exist within regional contexts. Having a regional perspective is also vital to recognize the relative value of an area for habitat or human use.

There are several regional organizations that may be interested in research projects similar to those presented in this SRA. As an example, the Northeast Regional Ocean Council (NROC), created in 2005 by the six New England governors, is responsible for facilitating the development and implementation of coordinated and collaborative regional goals and priorities to improve responses to regional issues and challenges. In 2011, NROC began a two-year work plan to include research initiatives for the waters surrounding the New England region related to:

- Characterizing baseline information on environmental, social and economic aspects of the region in order to inform coastal and marine spatial planning (CMSP) in the region;
- Identifying Areas for Potential Conservation in the region based on their ecological significance;
- Identifying areas in the region for future use, such as offshore renewable energy (ORE) development, or areas significant for commercial or recreational fishing;
- Develop a performance monitoring and evaluation system as part of the CMSP plan to inform plan adaptation over time; and
- Provide regional management governance structure and coordination mechanisms for integrated state, tribal and federal CMSP and decision-making.
- In addition, the following organizations may be potential collaborators or funders:
  - The Northeast Sea Grant Consortium (NESGC). The NESGC is partnership of all the Sea Grant organizations in the Northeast and have a common goal to support research and outreach related to regional ocean issues.
  - The Nature Conservancy (TNC). TNC has conducted much work at the regional scale, which could be compared to work done within the Ocean SAMP area, and each could be used to inform the other.
  - The United States Navy (USN). The USN has indicated interest as partners in acoustic and modeling studies.

Coordinating future Ocean SAMP research efforts when possible with regional organizations will result in a more accurate understanding of the area. Furthermore, because ORE will likely be developed across New England, collaborating on research focused on measuring the impact of development may help reduce project time and costs.

## **7. Conclusion**

The Ocean Special Area Management Plan (SAMP) process and resulting document has moved Rhode Island to the forefront of marine spatial planning (MSP) and offshore renewable

energy (ORE) development at local and national scales. Prior to the Ocean SAMP process, knowledge was limited about the natural resources and human uses of the waters off Rhode Island's coast. Today, much more is understood about the Ocean SAMP study area, in part due to the tremendous efforts conducted as part of this MSP process, including the of compilation and synthesis of existing information and collection of new data.

Since the completion of the Ocean SAMP, Rhode Island has made efforts to continue to overcome the scientific and management challenges related to MSP and ORE development. The development of this Science Research Agenda (SRA) is one example of this effort. The SRA identifies research and tasks that need to be accomplished in order to help guide scientifically sound management decisions, and was developed through a transparent and public process. Specifically, the SRA focuses on work to:

- Assist in obtaining and directing future funding opportunities;
- Continue to learn about and provide a better understanding of the current state of the Ocean SAMP ecosystem (environmental and biological conditions);
- Establish monitoring studies to assess changes in baseline conditions due to natural events and human activities;
- Investigate the potential effects of future offshore uses, including ORE development, and of global climate change;
- Advance the technology and efficiency of ORE devices;
- Develop an information framework to effectively communicate information;
- Facilitate collaborative, multidisciplinary research projects; and
- Support regional MSP efforts.

The SRA is also a mechanism to ensure the Ocean SAMP is implemented using an adaptive management approach. Since its inception in 1971, the Coastal Resource Management Council (CRMC) has managed Rhode Island's coastal waters using an adaptive management approach, a systematic process for continually improving management by adjusting practices and policies

as new information becomes available. This approach requires careful implementation, monitoring, evaluation of results, and adjustment of objectives and practices.

While this research agenda focuses on the Ocean SAMP study area, the research ideas expressed may be applicable for regional MSP efforts or valuable on a national scale as it relates to understanding the effects of ORE development.

## **8. References**

- Andrade, H. and Renaud, P.E., 2011. Polychaete/amphipod ratio as an indicator of environmental impact related to offshore oil and gas production along the Norwegian continental shelf. *Marine Pollution Bulletin*, 62(12): pp. 2836-44.
- Auster, P.J., Heinonen, K.B., Witharana, C., and McKee, M. 2009. A habitat classification scheme for the Long Island Sound region. Long Island Sound Study Technical Report. EPA Long Island Sound Office, Stamford, Connecticut 06904: 83 pp.
- Bonzek, C.F. J.Gartland, R.A. Johnson, and J.D. Lange Jr. 2008. NEAMAP Near Shore Trawl Survey: Peer Review Documentation. A report to the Atlantic States Marine Fisheries Commission. CEFAS 2011 (in press)  
[http://www.neamap.net/publications/VIMS\\_NEAMAP\\_Peer\\_Review\\_Documents.pdf](http://www.neamap.net/publications/VIMS_NEAMAP_Peer_Review_Documents.pdf)
- Bundesamt für Seeschifffahrt und Hydrographie [BSH]. 2007. *Standard Investigation of the Impacts of Offshore Wind Turbines on the Marine Environment* (StUK 3). BSH-Nr. 7003, Bundesamt für Seeschifffahrt und Hydrographie, Hamburg and Rostock, Germany. Available online from: <http://www.bsh.de/en/Products/Books/Standard/7003eng.pdf>
- CEFAS. 2004. Offshore Wind Farms: Guidance Note for Environmental Impact Assessment in Respect of FEPA and CPA Requirements.
- Crowder, L., and E. Norse. Essential ecological insights for marine ecosystem-based management and marine spatial planning. *Marine Policy*. 32: 772-778.
- Degraer, S., Brabant, R., and Rumes, B. (Eds.). 2011. Offshore wind farms in the Belgian part of the North Sea: Selected findings from the baseline and targeted monitoring. Royal Belgian Institute of Natural Sciences, Management Unit of the North Sea Mathematical

Models. Marine ecosystem management unit: 157 pp. + annex. Available online: [http://www.mumm.ac.be/Downloads/monwin\\_report\\_2011\\_final.pdf](http://www.mumm.ac.be/Downloads/monwin_report_2011_final.pdf) (Last accessed 4/4/2012).

Douvre, F. 2008. The engagement of stakeholders in the marine spatial planning process.

*Marine Policy*. 32: 762-771.

Durbin, E.G., Campbell, R.G., Casas, M.C., Ohman, M.D., Neihoff, B. Runge, J., and Wagner, M. 2003. Interannual variation in phytoplankton blooms and zooplankton productivity and abundance in the Gulf of Maine during winter. *Marine Ecology Progress Series*. 254: pp. 81-100.

Ehler, C. 2008. Conclusions: Benefits, lessons learned, and future challenges of marine spatial planning. *Marine Policy*. 32: 840-843.

Fisher, J., Frank, K.T., Kostylev, V.E., Shackell, N.L., Horsman, T., and Hannah, C.G. 2011. Evaluating a habitat template model's predictions of marine fish diversity on the Scotian Shelf and Bay of Fundy, Northwest Atlantic. *ICES Journal of Marine Science*, 68(10): pp. 2096-2105.

Freidland, K. D., Hare, J.A., Wood, G.B., Col, L.A., Buckley, L.J., Mountain, D.G., Kane, J., Brodziak, J., Lough, R.G. and Pilskaln, C.H. 2008. Does the fall phytoplankton bloom control recruitment of Georges Bank haddock, *Melanogrammus aeglefinus*, through parental condition? *Canadian Journal of Fisheries & Aquatic Science*. 65: pp. 1076-1086.

Fry, B. 1988. Food web structure on Georges Bank from stable C, N and S isotopic compositions. *Limnology and Oceanography*. 33 (5): pp. 1182-1190.

Fulweiler, R.W., Nixon, S.W., Buckley, B.A., and Granger, S.L. 2007. Reversal of the net nitrogen flux in coastal marine sediments. *Nature*. 448: pp. 180-182.

Fulweiler, R.W. and Nixon, S.W. 2009. Responses of benthic–pelagic coupling to climate change in a temperate estuary. *Eutrophication in Coastal Ecosystems*. 290: pp. 147-156.

Garrison, L.P. and Link, J.S. 2000. Dietary guild structure of the fish community in the Northeast United States continental shelf ecosystem. *Marine Ecology Progress Series*. 202: pp. 231-240.

- Global Insight. 2008. *Rhode Island Tourism: Strength in a Difficult Time: 2007 Tourism Satellite Account*.
- Grilli, A.R., Lado T., and Spaulding, M.L. 2012. A protocol to include ecosystem services in a wind farm cost model. *Journal of Environmental Engineering* (Under press/ online from June 29, 2012).
- Hansson, S., Hobbie, J.E., Elmgren, R., Larsson, U., Fry, B., and Johansson, S. 1997. The stable nitrogen isotope ratio as a marker of food-web interactions and fish migration. *Ecology*. 78 (7): pp. 2249-2257.
- Head, E.J.H., Brickman, D. and Harris, L.R. 2005. An exceptional haddock year class and unusual environmental conditions on the Scotian Shelf in 1999. *J. Plankton Res.* 27: pp. 597-602.
- Hyslop, E.J. 1980. Stomach contents analysis – a review of methods and their application. *Journal of Fish Biology*. 17 (4): pp. 411-429.
- Jolly, G.M., and Hampton, I. 1990. A stratified random transect design for acoustic surveys of fish stocks. *Canadian Journal of Fisheries and Aquatic Sciences*. 47 (7): pp. 1282-1291.
- LaFrance, M., Shumchenia, E., King, J., Pockalny, R., Oakley, B., Pratt, S., and Boothroyd, J. 2010. Benthic Habitat Distribution and Subsurface Geology Selected Sites from the Rhode Island Ocean Special Area Management Study Area. Technical Report 4. 99 pp; Kingston, RI, University of Rhode Island. Available online: [http://seagrant.gso.uri.edu/oceansamp/pdf/appendix/04-LaFrance\\_et\\_al\\_SAMPReport.pdf](http://seagrant.gso.uri.edu/oceansamp/pdf/appendix/04-LaFrance_et_al_SAMPReport.pdf) (Last accessed 3/10/2012).
- Love, M, J. Hyland, A. Ebeling, T. Herrlinger, A. Brooks, and E. Imamura. 1994. A pilot study of the distribution and abundances of rockfishes in relation to natural environmental factors and an offshore oil and gas production platform off the coast of Southern California. *Bulletin of Marine Science*, 53 (2-3): 1062-1085.
- Malek, A.J., LaFrance, M., Collie, J., and King, J. 2010. Fisheries ecology in Rhode Island and Block Island Sounds for the Rhode Island Ocean Special Area Management Plan 2010. Technical Report 14. 56 pp; Kingston, RI, University of Rhode Island. Available online:

[http://seagrant.gso.uri.edu/oceansamp/pdf/appendix/14-Maleketal\\_RISAMP.pdf](http://seagrant.gso.uri.edu/oceansamp/pdf/appendix/14-Maleketal_RISAMP.pdf)

(Last accessed 3/10/2012).

- Mather, I.R. and G. P. Watts, Jr. 1998. Geographic Information Systems for Cultural Resource Management and Site Specific Investigations. In: *Underwater Archaeology 1998*, edited by Lawrence E. Babits. Tucson, Arizona: Society for Historical Archaeology.
- Mather, I.R. and G. P. Watts, Jr. 2002. Geographic Information Systems. In: *International Handbook of Underwater Archaeology*, Carol V. Ruppé and Janet F. Barstad, editors. Plenum Publishing Corporation.
- Nixon, S.W., Fulweiler, R.W., Buckley, B.A., Granger, S.L., Nowicki, B.L. and Henry, K.W. 2009. The impact of changing climate on phenology, productivity, and benthic–pelagic coupling in Narragansett Bay. *Estuarine, Coastal and Shelf Science*. 82: pp. 1-18.
- Oumeraci, H., Jensen, J., Gönnert, G., Pasche, E., Kortenhaus, A., Naulin, M., Wahl, T., Thumm, S., Ujeyl, G., Gershovich, I., and Burzel, A. 2009. Flood risk analysis for a megacity: The German XtremRisK project. *Proc. Conference on Road Map towards a Flood Resilient Urban Environment*, Paris, France: pp. 8.
- Oviatt, C.A., Keller, A. and Reed, L. 2002. Annual Primary Production in Narragansett Bay with no Bay-Wide Winter–Spring Phytoplankton Bloom. *Estuarine, Coastal and Shelf Science*. 54: pp. 1013-1026.
- Oviatt, C.A. 2004. The Changing Ecology of Temperate Coastal Waters During a Warming Trend. *Estuaries*. 27(6): pp. 895-904.
- Pershing, A.J., Greene, C.H., Jossi, J.W., O'Brien, L., Brodziak, J.K.T. and Bailey, B. 2005. Interdecadal variability in the Gulf of Maine zooplankton community, with potential impacts on fish recruitment. *ICES Journal of Marine Science*. 62: pp. 1511-1523.
- Platt, T., Fuentes-Yaco, C. and Frank, K.T. 2003. Spring algal bloom and larval fish survival. *Nature*. 423: pp. 398-399.
- Pomeroy, R., and F. Douvre. 2008. The engagement of stakeholders in the marine spatial planning process. *Marine Policy*. 32: 816-822.

- Rademacher, K.R. and J.H. Render. 2003. Fish assemblages around oil and gas platforms in the Northeastern Gulf of Mexico: Developing a survey design. In: *Fisheries, Reefs, and Offshore Development*. Stanley, D.R. and Scarborough-Bull, A., eds. American Fisheries Society Symposium 36. Bethesda, Maryland: American Fisheries Society.
- Rhetore, P.E., Fuglsang, P., Larsen, G., Buhl, T., Larsen, T.J. and Madsen, H.A. 2011. TopFarm: Multi-fidelity optimization of offshore wind farm. In *Proc. 20th Intl. Offshore and Polar Engng. Conf.* (Maui, Hawaii, USA, June 19-24).
- Rice, J., Arvanitidis, C., Borja, A., Frid, C., Hiddink, J.G., and Krause, J. 2012. Indicators for Sea-floor Integrity under the European Marine Strategy Framework Directive. *Ecological Indicators*, 12(1): pp. 174-184.
- Rhode Island Coastal Resources Management Council (RI CRMC). 2010. Rhode Island Ocean Special Area Management Plan. Adopted October 19, 2010. Kingston, RI, University of Rhode Island. Available online: <http://seagrant.gso.uri.edu/oceansamp/documents.html> (Last accessed 5/21/12).
- Saunders, G., Bedford, G.S., Trendall, J.R., and Sotheran, I. (2011). Guidance on survey and monitoring in relation to marine renewables deployments in Scotland. Volume 5. Benthic Habitats. Unpublished draft report to Scottish Natural Heritage and Marine Scotland
- Smith, L.M., Whitehouse, S. and Oviatt, C.A. 2010. Impacts of climate change on Narragansett Bay. *Northeast Naturalist*. 17(1): pp. 77-90.
- Spaulding, M.L., Grilli, A.R., Damon, C., and Fugate, G. 2010. Application of Technology Development Index and principal component analysis and cluster methods to ocean renewable energy facility siting. *Marine Technology Society Journal*. 44(1): pp. 8-23.
- St. Martin, K., and M. Hall-Arber. 2008. The missing layer: Geo-technologies, communities, and implications for marine spatial planning. *Marine Policy*. 32: 779-786.
- Steele, J.H. 2001. Network analysis of food webs: pp. 1870-1875.
- Thunberg, E. 2008. Trends in Selected Northeast Region Marine Industries. Woods Hole, MA: National Oceanic and Atmospheric Administration Northeast Fisheries Science Center, July 2008.

- Tyrrell, T. J. and Johnston, R.J. 2001. A framework for assessing direct economic impacts of tourist events: distinguishing origins, destinations, and causes of expenditures. *Journal of Travel Research*, 40: pp. 94-100.
- Van den Eynde, D., Brabant, R., Fettweis, M., Francken, F., Melotte, J., Sas, M., Lancker, V.V., 2010. Monitoring of hydrodynamic and morphological changes at the C-Power and Belwind offshore windfarm sites – A synthesis, in: Degraer, S., Brabant, R., Rumes, B., 2010. Offshore wind farms in the Belgian part of the North Sea: Early environmental impact assessment and spatio-temporal variability. pp. 19-36.
- Verdant Power. 2010. *Roosevelt Island Tidal Energy Project, FERC No. 12611, Final Kinetic Hydropower Pilot Licence Application, Volume 4, Part 1: Proposed Monitoring Plans, RITE Monitoring of Environmental Effects (RMEE)*. Verdant Energy, LLC, New York, NY.D144+D167 <http://www.theriteproject.com/uploads/VPRITE-FLAVol4-Part1of3.pdf>
- Verfaillie, E., Degraer, S., Schelfaut, K., Willems, W., and Van Lancker, V., 2009. A protocol for classifying ecologically relevant marine zones, a statistical approach. *Estuarine, Coastal and Shelf Science*, 83(2): pp. 175-185.
- Watts, Jr., G.P., and I.R. Mather. 1996. GIS for Historical and Cultural Resource Management in the Waters around Charleston. Data sets and report submitted to the City of Charleston, South Carolina.
- Watts, Jr., G.P., and I.R. Mather. 1997. Historic Properties Treatment Plan and Geographic Information System for the James River Navigation Project.” Data sets and report submitted to US Corps of Engineers, Norfolk District, Virginia.
- Wiley, D.N., Moller, J.C, and Zilinskas, K.A. 2003. The distribution and density of commercial fisheries and baleen whales within the Stellwagen Bank National Marine Sanctuary: July 2001-June 2002. *MTS Journal* 37 (1): 35-53.
- Wilhelmsson, D., Malm, T., and Ohman, M. 2006. The influence of offshore windpower on demersal fish. *ICES Journal of Marine Science*, 63: 775-78.
- Winiarski, K.J., Paton, P., Trocki, C.L., and McWilliams, S.R. 2011. Spatial distribution, abundance, and flight ecology of birds in nearshore and offshore waters of Rhode Island:

January 2009 to August 2010. Rhode Island Ocean Special Area Management Plan  
Interim Report.

**Appendix: Monitoring Protocols Developed by URI through the National Oceanographic Partnership Program (NOPP)**

**\*\*Full NOPP report available here:**

<http://www.data.boem.gov/PI/PDFImages/ESPIS/5/5208.pdf>

The tables below are summaries of the NOPP monitoring protocols and examples of methods to accomplish them. For these tables “cost” is described as low (<\$100,000), moderate (\$100,000-\$500,000), and high (>\$500,000) per year. For more information, the reader to referred to the full NOPP report.

**Benthic Habitat and Ecology**

**Benthic Environment Monitoring Protocol 1: Sediment Scour and/or Deposition.** Assess changes in sediment scour and/or deposition around ORE devices.

<b>Indicator(s) of the impact</b>	Scour: increase in median grain size; decrease in organic content; decrease in seabed volume  Deposition: decrease in median grain size; increase in organic content; increase in seabed volume	
<b>Methodology or Technique to Collect Data</b>	Particle size analysis; Multibeam/interferometric bathymetry	
<b>Description of Methodology or Technique(s) for collecting data</b>	Seasonal surveys, 5 years  Grain size: *5-sample transect at 3 devices out to 200m  Bathymetry: overlapping transects for 100% coverage (at least 0.5 m pixels) 1 km radius at 3 devices	Annual surveys, 3 years  Grain size: *3-sample transect at 3 devices out to 200m  Bathymetry: overlapping transects for 100% coverage (at least 0.5 m pixels) 500m radius 3 devices
<b>Methodology for Analyzing data</b>	ANOVA on median grain size; Volume change estimate using mosaicked bathymetry models	

<b>Frequency and Duration</b>	1 preconstruction survey; Seasonal operation; 1 post-construction survey	1 preconstruction survey; Annual operation; 1 post-construction survey
<b>Spatial Scale</b>	200m – 1km radius around 3 devices	500m radius around 3 devices
<b>How well does methodology account for environmental variability?</b>	Seasonal and interannual variability	Interannual variability
<b>Cost</b>	Moderate	Low
<b>Relationship to Other Protocols:</b>	Can be combined with benthic community composition monitoring protocol (Protocol Z2)	
<b>Data Format:</b>	Data table time series	
<b>Data Output:</b>	Time series values for median grain size and standard deviations; time series on volume at each turbine and standard deviation	
<b>Examples where methodology has been used:</b>	Degraer et al., 2011; Saunders et al., 2001	

**Benthic Environment Monitoring Protocol 2: Changes in benthic community composition.**

Assess changes in benthic community composition at or in vicinity of ORE installation.

<b>Indicator(s) of the impact</b>	Change in abundance, diversity, % cover, multivariate community composition	
<b>Methodology or Technique to Collect Data</b>	Grab samples (Smith McIntyre or similar) ~0.1m <sup>2</sup> /sample (soft bottom); Underwater video transects (soft and hard bottom)	
<b>Description of Methodology or Technique(s) for collecting data</b>	Seasonal surveys, 5 years *5-sample transect at 5 devices out to 200m **200m UWvideo at each device AND at reference station <1km	Annual surveys, 3 years *3-sample transect at 3 devices out to 200m **200m UWvideo at 3 device AND at reference station <1km
<b>Methodology for Analyzing data</b>	ANOVA on abundance, diversity, % cover; ANOSIM on community composition: over time and between ORED and reference	
<b>Frequency and Duration</b>	1 preconstruction survey; Seasonal	1 preconstruction survey; Annual operation; 1 post-

	operation; 1 post-construction survey	construction survey
<b>Spatial Scale</b>	200 m radius around 5 devices	200m radius around 3 devices
<b>How well does this methodology account for environmental variability?</b>	Seasonal and interannual variability	Interannual variability
<b>Cost</b>	Moderate	Low
<b>Relationship to Other Protocols:</b>	Can be combined with scour/deposition monitoring protocol (Protocol Z1) Can be combined with reef effects monitoring protocol (Protocol Z3)	
<b>Data Format:</b>	Data table time series	
<b>Data Output:</b>	Time series values for abundance, diversity, % cover and standard deviations; time series summary metric on benthic community composition (Indicator species, SIMPER, etc.)	
<b>Examples where this methodology has been used:</b>	Degraer et al., 2011; Saunders et al., 2001	

**Benthic Environment Monitoring Protocol 3: Reef Effects.** Assess change (e.g. increase) in hard bottom habitat (reef effect) and non-native species at or in vicinity of ORE installation.

<b>Indicator(s) of the impact</b>	Increase in % cover, biomass of epifaunal organisms; increase in presence of non-native species	
<b>Methodology or Technique to Collect Data</b>	Diver imagery and scrape samples	
<b>Description of Methodology or Technique(s) for collecting data</b>	Seasonal surveys, 5 years  Diver picture/video, then scrape and collect 0.25x0.25 m quadrat; 3 quadrats per device (high, med low water); 3 devices	Annual surveys, 3 years  3 ROV video transect per device; 3 devices
<b>Methodology for Analyzing data</b>	% cover estimate from imagery, dry weight biomass; ANOVA	ANOVA on % cover, # species

<b>Frequency and Duration</b>	Seasonal during operation only	Annual during operation only
<b>Spatial Scale</b>	Small	Large
<b>How well does this methodology account for environmental variability?</b>	Seasonal and interannual variability	Interannual variability
<b>Cost</b>	Moderate	Low
<b>Relationship to Other Protocols:</b>	Can be combined with Fisheries Monitoring Protocol X3	
<b>Data Format:</b>	Data table time series	
<b>Data Output:</b>	time series values for % cover, biomass; presence absence of non-native species	
<b>Examples where this methodology has been used:</b>	Degraer et al., 2011	

**Benthic Environment Monitoring Protocol 4: Hydrodynamics.** Assess changes in hydrodynamics due to ORE development.

<b>Indicator(s) of the impact</b>	Change in residual flow rates; change in water column turbidity	
<b>Methodology or Technique to Collect Data</b>	Bottom-mounted acoustic Doppler current profilers (ADCPs); Turbidity sensors	
<b>Description of Methodology or Technique(s) for collecting data</b>	Seasonal surveys, 5 years  3 ADCP/turbidity sensor package in ORED; 1 ADCP/turbidity at reference site	Annual surveys, 3 years  1 ADCP/turbidity sensor package in ORED; 1 ADCP/turbidity at reference site
<b>Methodology for Analyzing data</b>	ANOVA time-average flow velocity; time-average turbidity	
<b>Frequency and Duration</b>	Preconstruction baseline survey; seasonal averages during operation	Preconstruction baseline survey; Annual averages during operation
<b>Spatial Scale</b>	Transect across entire	Point location within

	development	development
<b>How well does this methodology account for environmental variability?</b>	Depending on length of deployment, captures from tidal to interannual variability	
<b>Cost</b>	Moderate	Low
<b>Data Format:</b>	Data table time series	
<b>Data Output:</b>	time series for time-averaged flow rates and turbidity values (tidal frequency, daily, monthly, seasonally, annually)	
<b>Examples where this methodology has been used:</b>	Van den Eynde et al., 201	

**Avian**

**Avian Protocol 1: Ship-based Visual Surveys.** Assess changes in spatial distribution and abundance of marine birds\* due to ORE development. (*Can also be applied for pre-siting baseline studies.*)

<b>Indicator(s) of Impact</b>	Spatially-explicit changes in density estimates.
<b>Methodology or Technique to Collect Data</b>	Ship-based line-transect surveys using at least two observers in a 20-100 m ship.
<b>Description of Methodology or Technique(s) for Collecting Data</b>	Line-transect distance sample technique (Camphuysen et al. 2004; See Appendix A).
<b>Methodology for Analyzing data</b>	<ul style="list-style-type: none"> <li>• Before After Gradient (BAG) or Before After Control Impact (BACI) monitoring design.</li> <li>• Model based analysis (See Petersen <i>et al.</i> 2011).</li> </ul>
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>• Minimum of three surveys per season (winter, spring, summer, and fall) to monitor different migratory species.</li> <li>• Baseline: Minimum of two years pre construction (could be &lt;2 years if adequate historical baseline data exists).</li> <li>• Post-construction: Recommended for up to 15 years post construction in years 1, 2, 3, 5, 10 and 15.</li> </ul>

<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>• Sites &lt;5 km<sup>2</sup>, a buffer<sup>#</sup> of at least 1 km around impact area.</li> <li>• Sites 5 km<sup>2</sup> – 10 km<sup>2</sup>, a buffer of at least 2 km.</li> <li>• Sites &gt;10 km<sup>2</sup>, a buffer of at least 4 km.</li> <li>• <sup>#</sup>Not necessarily a symmetrical buffer depending on ORED device and predicted environmental effects.</li> </ul>
<b>Strengths</b>	<ul style="list-style-type: none"> <li>• High detection probabilities for most species of marine birds.</li> <li>• Possible to collect other covariates simultaneously including: sea surface temperature, chlorophyll, salinity and bioacoustics data.</li> <li>• Ability to identify individuals to species for targeted taxa (e.g., federally-listed Roseate Terns).</li> <li>• Ability to determine age and/or gender for some species to model population dynamics.</li> <li>• Surveys prior to 2005 were ship-based surveys, so this method is directly comparable to archived data.</li> <li>• Can be conducted nearshore (&lt;3miles from shore), assuming appropriate water depth.</li> <li>• Safety risk for crew members lower than aerial surveys.</li> <li>• Able to potentially survey other taxa including marine mammals and sea turtles simultaneously.</li> </ul>
<b>Limitations</b>	<ul style="list-style-type: none"> <li>• Not suitable for nocturnal migrants including many species of shorebirds (Charadriiformes) and songbirds (Passeriformes).</li> <li>• Slow survey speed, thus relatively small areas can be surveyed within a day.</li> <li>• Sea state limitations (especially in winter when favorable conditions are limited) Not suitable for some disturbance-prone species (e.g., seaducks, loons).</li> <li>• May displace or attract species, resulting in biased density estimates.</li> <li>• Ships may not be allowed within some wind facilities making post-construction comparisons difficult.</li> <li>• Cannot be conducted in very shallow or rocky areas.</li> <li>• Glare/wind/waves can often affect detection probabilities.</li> <li>• Challenging in areas with strong tidal currents.</li> </ul>
<b>Relationship to Other Protocols</b>	<ul style="list-style-type: none"> <li>• Can be conducted in combination with Marine Mammals and Sea Turtles Protocol W1 and Fisheries Resources Protocol X1</li> </ul>
<b>Cost</b>	<ul style="list-style-type: none"> <li>• Moderate (depends on number of surveys)</li> </ul>
<b>Type of Data Collected</b>	<ul style="list-style-type: none"> <li>• Spatially-explicit locations with an accuracy of ± 300 m.</li> <li>• Flock size and identification of most individuals to species.</li> <li>• Distance of individual or flock from transect centerline to model density estimates.</li> <li>• Behavior (foraging, resting based on location – either on the water or flying).</li> </ul>

	<ul style="list-style-type: none"> <li>• For birds in flight, flight direction and estimated flight altitude.</li> <li>• Environmental covariates (sea state, wind speed, wind direction).</li> </ul>
<b>Data Output</b>	<ul style="list-style-type: none"> <li>• Spatially-explicit density estimates (and associated variance) by species within and outside the development area.</li> </ul>
<b>Examples where this technique has been used for ORED monitoring</b>	<ul style="list-style-type: none"> <li>• Vanermen et al. 2010</li> </ul>

\*Marine birds suitable for these survey include loons (Gaviidae), grebes (Podicipedidae), tubenoses (Procellariidae, Hydrobatidae), pelicans and allies (Fregatidae, Pelecanidae, Phaethontidae, Sulidae), cormorants (Phalacrocoracidae), wading birds (Ardeidae), waterfowl (Anatidae), diurnal raptors (Accipitridae, Falconidae), phalaropes (Phalaropus spp.), gulls and allies (Laridae), terns (Sterna spp.), and alcids (Alcidae).

**Avian Protocol 2: Aerial Visual Surveys using Human Observers.** Assess changes in spatial distribution and abundance of marine birds\* due to ORE development. (Can also be applied for pre-siting baseline studies)

<b>Indicator(s) of Impact</b>	Spatially-explicit changes in density estimates.
<b>Methodology or Technique to Collect Data</b>	Aerial line-transect; Visual surveys by observers.
<b>Description of Methodology or Technique(s) for Collecting Data</b>	Line-transect Distance Sample Technique (Camphuysen et al. 2004; Appendix A) from a plane with at least two observers.
<b>Methodology for Analyzing data</b>	<ul style="list-style-type: none"> <li>• Before After Gradient (BAG) or Before After Control Impact (BACI) monitoring design.</li> <li>• Model based analysis (See Petersen <i>et al.</i> 2011).</li> </ul>
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>• Minimum of three surveys per season (winter, spring, summer, and fall) to monitor different migratory species.</li> <li>• Baseline: Minimum of two years pre construction (could be &lt;2 years if adequate historical baseline data exists).</li> <li>• Post-construction: Recommended for up to 15 years post construction in years 1, 2, 3, 5, 10 and 15.</li> </ul>
<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>• Sites &lt;5 km<sup>2</sup>, a buffer of at least 1 km around impact area.</li> <li>• Sites 5 km<sup>2</sup> – 10 km<sup>2</sup>, a buffer of at least 2 km.</li> <li>• Sites &gt;10 km<sup>2</sup>, a buffer of at least 4 km.</li> <li>• *Not necessarily a symmetrical buffer depending on device and predicted environmental effects.</li> </ul>
<b>Strengths</b>	<ul style="list-style-type: none"> <li>• High detection probabilities for most species of marine birds.</li> <li>• Able to detect disturbance-prone species.</li> <li>• Able to potentially simultaneously survey marine mammals.</li> </ul>
<b>Limitations</b>	<ul style="list-style-type: none"> <li>• Not suitable for nocturnal migrants including many species of shorebirds (Charadriiformes) and songbirds (Passeriformes).</li> <li>• Birds disturbed due to low flight altitude (generally after being recorded).</li> <li>• Some detections may be identified only to a species group.             <ul style="list-style-type: none"> <li>• Found to underestimate abundance of cryptic species.</li> </ul> </li> <li>• Some wind facilities may not allow flights post-construction.</li> <li>• Safety issue for low altitude flights compared to ship-based surveys.</li> <li>• Glare can often affect detection probabilities.</li> </ul>

<b>Relationship to Other Protocols</b>	<ul style="list-style-type: none"> <li>• Can be conducted in combination with Marine Mammals &amp; Sea Turtles Protocol W1, and Fisheries Resources Protocol X1</li> </ul>
<b>Cost</b>	<ul style="list-style-type: none"> <li>• Moderate (depends on number of surveys)</li> </ul>
<b>Type of Data Collected</b>	<ul style="list-style-type: none"> <li>• Spatially-explicit locations with an accuracy of <math>\pm 100</math> m.</li> <li>• Flock size and identification of some detections to species or species groups.                             <ul style="list-style-type: none"> <li>• Distance of individual or flock from transect centerline to model density estimates.</li> <li>• Behavior (foraging, resting based on location - on water or flying).</li> <li>• For birds in flight, flight direction and estimated flight altitude.</li> </ul> </li> <li>• Environmental covariates (sea state, wind speed, wind direction).</li> </ul>
<b>Data Output</b>	<ul style="list-style-type: none"> <li>• Spatially-explicit density estimates (and associated variance) by species / taxonomic groups within and outside development area.</li> </ul>
<b>Examples where this technique has been used for ORED monitoring</b>	Petersen et al. 2011; Maclean et al. 2006

\*Marine birds suitable for these survey include loons (Gaviidae), tubenoses (Procellariidae, Hydrobatidae), pelicans and allies (Fregatidae, Pelecanidae, Phaethontidae, Sulidae), cormorants (Phalacrocoracidae), wading birds (Ardeidae), waterfowl (Anatidae), gulls and allies (Laridae), terns (Sterna spp.), and alcids (Alcidae).

**Avian Protocol 3: Aerial Surveys using High Definition Videography.** Assess changes in spatial distribution and abundance of marine birds\* due to ORE development. *(Can also be applied for pre-siting baseline studies)*

<b>Indicator(s) of Impact</b>	Spatially-explicit changes in density estimates.
<b>Methodology or Technique to Collect Data</b>	Aerial strip-transect surveys; High definition videography.
<b>Description of Methodology or Technique(s) for Collecting Data</b>	Strip transect methodology (See Appendix A).
<b>Methodology for Analyzing data</b>	<ul style="list-style-type: none"> <li>• Before After Gradient Before After Control Impact (BACI) monitoring design.</li> <li>• Model based analysis (See Petersen et al. 2011).</li> </ul>

<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>• Minimum of three surveys seasonally.</li> <li>• Baseline: At least two years pre construction (maybe possible in &lt;2 years if adequate historical baseline data exists). If year 1 and year 2 are very different years for certain particular common and abundant species than baseline surveys should be continued.</li> <li>• Post-construction: Recommended for up to 15 years post construction in years 1, 2, 3, 5, 10 and 15.</li> </ul>
<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>• Sites &lt;5 km<sup>2</sup>, a buffer of at least 1 km around impact area.</li> <li>• Sites 5 km<sup>2</sup> – 10 km<sup>2</sup>, a buffer of at least 2 km.</li> <li>• Sites &gt;10 km<sup>2</sup>, a buffer of at least 4 km.</li> <li>• *Not necessarily a symmetrical buffer depending on device and predicted environmental effects.</li> </ul>
<b>Strengths</b>	<ul style="list-style-type: none"> <li>• High detection probabilities for all marine birds</li> <li>• Large area can be surveyed rapidly.</li> <li>• Flight elevation high enough to not disturb birds, thus able to sample disturbance prone species.</li> <li>• Safer than observer-based aerial surveys that fly at lower altitudes.</li> <li>• More aircraft are potentially suitable for videography, as they do not have to be high winged aircraft.</li> <li>• Permanent record of observations that could be reviewed by biologists in the future.</li> <li>• Spatially-explicit density estimates.</li> <li>• Potential to estimate of flight altitude of individuals or flocks.</li> <li>• Able to survey marine mammals and sea turtles simultaneously.</li> </ul>
<b>Limitations</b>	<ul style="list-style-type: none"> <li>• Not suitable for nocturnal migrants such shorebirds (Charadriiformes) and songbirds (Passeriformes).</li> <li>• Technology is still evolving.</li> <li>• Similar species may not be identifiable depending on imagery.</li> </ul>
<b>Relationship to Other Protocols</b>	<ul style="list-style-type: none"> <li>• Can be conducted in combination with Marine Mammals &amp; Sea Turtles Protocol W1, and Fisheries Resources Protocol X1</li> </ul>
<b>Cost</b>	<ul style="list-style-type: none"> <li>• High (depends on number of surveys)</li> </ul>
<b>Type of Data Collected</b>	<ul style="list-style-type: none"> <li>• Spatially-explicit density estimates of individuals or flocks to within 100 m of actual locations.</li> <li>• Flock size and identification of some detections to species or species groups (e.g. alcids).             <ul style="list-style-type: none"> <li>• Distance of individual or flock from transect centerline to model density estimates.</li> <li>• Behavior (foraging, resting based on location - on water or flying).</li> </ul> </li> <li>• Environmental covariates (sea state, wind speed, wind direction).</li> </ul>

<b>Data output</b>	<ul style="list-style-type: none"> <li>Spatially-explicit density estimates (and associated variance) by species / taxonomic groups within and outside development area.</li> </ul>
<b>Examples where this technique has been used for ORED monitoring</b>	Mellor et al. 2007; Mellor and Maher 2008; Buckland et al. 2012

\*Marine birds suitable for these survey include loons (Gaviidae), grebes (Podicipedidae), tubenoses (Procellariidae, Hydrobatidae), pelicans and allies (Fregatidae, Pelecanidae, Phaethontidae, Sulidae), cormorants (Phalacrocoracidae), wading birds (Ardeidae), waterfowl (Anatidae), diurnal raptors (Accipitridae, Falconidae), phalaropes (Phalaropus spp.), gulls and allies (Laridae), terns (Sterna spp.), and alcids (Alcidae).

**Avian Protocol 4: Aerial Surveys using Digital Still Photography.** Assess changes in spatial distribution and abundance of marine birds\* due to ORE development. *(Can also be applied for pre-siting baseline studies)*

<b>Indicator(s) of Impact</b>	Spatially-explicit changes in density estimates.
<b>Methodology or Technique to Collect Data</b>	Aerial strip-transect surveys: Digital still photography.
<b>Description of Methodology or Technique(s) for Collecting Data</b>	Strip transect methodology (See Appendix A).
<b>Methodology for Analyzing data</b>	<ul style="list-style-type: none"> <li>Before After Gradient monitoring design.</li> <li>Model based analysis (See Petersen et al. 2011).</li> </ul>
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>Minimum of three surveys seasonally.</li> <li>Baseline: At least two years pre construction (maybe possible in &lt;2 years if adequate historical baseline data exists) . If year 1 and year 2 are very different years for certain particular common and abundant species than baseline surveys should be continued.</li> <li>Post-construction: Recommended for up to 15 years post construction in years 1, 2, 3, 5, 10 and 15.</li> </ul>
<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>Sites &lt;5 km<sup>2</sup>, a buffer of at least 1 km around impact area.</li> <li>Sites 5 km<sup>2</sup> – 10 km<sup>2</sup>, a buffer of at least 2 km.</li> <li>Sites &gt;10 km<sup>2</sup>, a buffer of at least 4 km.</li> <li>*Not necessarily a symmetrical buffer depending on device and predicted environmental effects.</li> </ul>

<b>Strengths</b>	<ul style="list-style-type: none"> <li>• High detection probabilities for most marine birds depending on image quality.</li> <li>• Large study areas can be surveyed rapidly.</li> <li>• Due to high flight altitude, able to survey disturbance-prone species.</li> <li>• Safer than observer-based aerial surveys that fly at lower altitudes.</li> <li>• More aircraft are potentially suitable for videography, as they do not have to be high winged aircraft.</li> <li>• Permanent record of observations that could be reviewed by biologists in the future.</li> <li>• Spatially-explicit density estimates</li> <li>• Potentially estimate of flight altitude of individuals or flocks.</li> <li>• Able to survey marine mammals and sea turtles simultaneously.</li> </ul>
<b>Limitations</b>	<ul style="list-style-type: none"> <li>• Not suitable for nocturnal migrants such as plovers, sandpipers, and songbirds (Passeriformes).</li> <li>• Technology is still evolving.</li> <li>• Similar species may not be identifiable.</li> </ul>
<b>Relationship to Other Protocols</b>	<ul style="list-style-type: none"> <li>• Can be conducted in combination with Marine Mammals &amp; Sea Turtles Protocol W1, and Fisheries Resources Protocol X1</li> </ul>
<b>Cost</b>	<ul style="list-style-type: none"> <li>• High (depends on number of surveys)</li> </ul>
<b>Type of Data Collected</b>	<ul style="list-style-type: none"> <li>• Spatially-explicit density estimates of individuals or flocks to within 100 m of actual locations.</li> <li>• Flock size and identification of some detections to species or species groups.                             <ul style="list-style-type: none"> <li>• Distance of individual or flock from transect centerline to model density estimates.</li> <li>• Behavior (foraging, resting based on location - on water or flying).</li> </ul> </li> <li>• Environmental covariates (sea state, wind speed, wind direction).</li> </ul>
<b>Data output</b>	<ul style="list-style-type: none"> <li>• Spatially-explicit density estimates (and associated variance) by species / taxonomic groups within and outside development area.</li> </ul>
<b>Examples where this technique has been used for ORED monitoring</b>	Buckland et al. 2012

\*Marine birds suitable for these survey include loons (Gaviidae), grebes (Podicipedidae), tubenoses (Procellariidae, Hydrobatidae), pelicans and allies (Fregatidae, Pelecanidae, Phaethontidae, Sulidae), cormorants (Phalacrocoracidae), wading birds (Ardeidae), waterfowl (Anatidae), diurnal raptors (Accipitridae, Falconidae), phalaropes (Phalaropus spp.), gulls and allies (Laridae), terns (Sterna spp.), and alcids (Alcidae).

**Avian Protocol 5: Radar Surveys.** Assess changes in avian movement ecology, such as migration and foraging flight paths between foraging and roosting sites due to ORE facility.

<b>Indicator(s) of Impact</b>	Changes in flight paths of foraging or commuting birds within development area.
<b>Methodology or Technique to Collect Data</b>	Marine Radar Surveys.
<b>Description of Methodology or Technique(s) for Collecting Data</b>	<ul style="list-style-type: none"> <li>• Marine Radar. X-band for vertical radar and either X-Band or S-band for horizontal radar.</li> <li>• Minimum of 25kW recommended and a vertical beam width of 20 degrees to 25 degrees and a horizontal beam of 0.9 degrees and a transmission frequency of about 9.4GHz (x-band radar).</li> <li>• Standard operating range should be 1.5 km for vertical and 3 km for horizontal radar.</li> <li>• Sea state &lt;5.</li> </ul>
<b>Methodology for Analyzing data</b>	<ul style="list-style-type: none"> <li>• A Before-After design would provide information on changes in movement patterns (Desholm and Kahlert 2005)</li> </ul>
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>• 1-2 years pre-construction/1-2 years post construction.</li> <li>• Continuous 24-hour monitoring</li> </ul>
<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>• Sites &lt;5 km<sup>2</sup>, a buffer of at least 1 km around impact area.</li> <li>• Sites 5 km<sup>2</sup> – 10 km<sup>2</sup>, a buffer of at least 2 km.</li> </ul>
<b>Strengths</b>	<ul style="list-style-type: none"> <li>• Flight trajectories can be stored in GIS.</li> <li>• Provides quantitative data on diurnal and nocturnal movements</li> <li>• Quantitative estimates of number of targets passing an area.</li> <li>• Flight speed can be used to group echoes to differentiate groups of birds (Smaller birds fly slower than larger birds).</li> <li>• Weather covariates can be collected simultaneously to investigate relationships with migration patterns.</li> <li>• Quantitative, accurate flight altitude information. Can be combined with ground truthing to make detections species specific.</li> </ul>
<b>Limitations</b>	<ul style="list-style-type: none"> <li>• Generally cannot identify targets to species.</li> <li>• Algorithms to analyze raw radar data are often proprietary and not directly comparable among studies.</li> <li>• For offshore developments, requires a stable platform for the radar unit, which can be challenging.</li> <li>• Wave and sea clutter can often make data unusable.</li> <li>• X-band radar more susceptible to rain clutter.</li> <li>• Can be challenging to develop precise quantitative counts due to issues with detection probabilities.</li> </ul>

<b>Cost</b>	<ul style="list-style-type: none"> <li>• High (due to need for stable platform)</li> </ul>
<b>Type of Data Collected</b>	<ul style="list-style-type: none"> <li>• Number of targets per hr by area and specific travel routes (3D).</li> </ul>
<b>Data Output</b>	<ul style="list-style-type: none"> <li>• Altitude distributions (100m increments up to 1000m).</li> <li>• Map of radar tracks (pre and post construction). For ground-truthed data, could provide some species-specific information.</li> <li>• Phenology of movements (number of targets, flight directions, and flight altitude).</li> </ul>
<b>Examples where this technique has been used for ORED monitoring</b>	Krijgsveld et al. 2010; Desholm et al. 2004.

**Avian Protocol 6: Visual Surveys of Flight Ecology.** Assess changes in avian movement ecology, such as migration and foraging flight paths between foraging and roosting sites due to ORE facility.

<b>Indicator(s) of Impact</b>	Changes in flight paths of foraging or commuting birds within development area.
<b>Methodology or Technique to Collect Data</b>	Visual Surveys (Ground-truthing radar surveys).
<b>Description of Methodology or Technique(s) for Collecting Data</b>	<ul style="list-style-type: none"> <li>• Visual observations to identify species detected by radar.</li> <li>• Observations conducted from a stable platform near radar unit where targets can be seen prior to entering buffer surrounding wind farm.</li> <li>• Communication recommended between observer and radar operator.</li> </ul>
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>• Surveys should take place during peak migration periods of target species.</li> <li>• This study could be conducted either pre- or post-construction (see Desholm and Kahlert 2005).</li> <li>• 1-2 years pre construction/1-2 years post construction?</li> </ul>
<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>• Depends on if X- or S-band radar is used, but generally within 1-3 km of radar unit.</li> </ul>
<b>Strengths</b>	<ul style="list-style-type: none"> <li>• Allows identification of radar targets to species level, but only for larger diurnal migrants.</li> <li>• Collected data could potentially be used to assess changes in flight ecology of target species following construction.</li> </ul>

<b>Limitations</b>	<ul style="list-style-type: none"> <li>• Not feasible for nocturnal targets or other low visibility conditions Detection probabilities are uncertain, but vary by size of targets Most useful for larger species (crow sized and larger).</li> <li>• Working on offshore platforms can be dangerous.</li> </ul>
<b>Cost</b>	<ul style="list-style-type: none"> <li>• Low for observers in coastal sites</li> </ul>
<b>Type of Data Collected</b>	<ul style="list-style-type: none"> <li>• Speciation of flocks or individuals recorded on radar.</li> </ul>
<b>Data Output</b>	<ul style="list-style-type: none"> <li>• Flight intensities (i.e., targets per hour) and flight altitude of target species.</li> </ul>
<b>Examples where this technique has been used for ORED monitoring</b>	<ul style="list-style-type: none"> <li>• Krijgsveld et al. 2010</li> </ul>

**Avian Protocol 7: Flight Call Surveys.** Assess changes in avian movement ecology, such as migration and foraging flight paths between foraging and roosting sites due to ORE facility.

<b>Indicator(s) of Impact</b>	Changes in flight paths of foraging or commuting birds within development area.
<b>Methodology or Technique to Collect Data</b>	Flight Call Surveys (Ground truth radar surveys).
<b>Description of Methodology or Technique(s) for Collecting Data</b>	<ul style="list-style-type: none"> <li>• Acoustic observations of flight calls to determine species composition of birds detected by radar during nocturnal surveys.</li> <li>• Observations conducted from a stable platform near radar unit where individuals or flocks can be heard prior to entering buffer surrounding wind farm.</li> <li>• Communication between observer and radar operator.</li> </ul>
<b>Methodology for Analyzing data</b>	<ul style="list-style-type: none"> <li>• Before – After – Control Impact.</li> </ul>
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>• Surveys should take place during peak migration periods of target species.</li> <li>• At least one year pre-construction and one year post construction.</li> </ul>
<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>• Development area.</li> </ul>
<b>Strengths</b>	<ul style="list-style-type: none"> <li>• Only way to identify radar targets to species at night.</li> <li>• Primarily useful for passerines.</li> </ul>
<b>Limitations</b>	<ul style="list-style-type: none"> <li>• Limited detection probability for calling targets. Some species do not call when flying at night.</li> <li>• Working on offshore platforms can be dangerous.</li> </ul>

<b>Cost</b>	<ul style="list-style-type: none"> <li>Moderate (due to working on offshore platform)</li> </ul>
<b>Type of Data Collected</b>	<ul style="list-style-type: none"> <li>Identification of radar targets to species.</li> </ul>
<b>Data Output</b>	<ul style="list-style-type: none"> <li>Relative flight call intensities (i.e., calls per hour by species).</li> </ul>
<b>Examples where this technique has been used for ORED monitoring</b>	Krijgsveld et al. 2010

**Avian Protocol 8: Radio Tracking.** Assess changes in avian movement ecology, such as migration and foraging flight paths between foraging and roosting sites due to ORE facility.

<b>Indicator(s) of Impact</b>	Changes in flight paths of foraging or commuting birds within development area.
<b>Methodology or Technique to Collect Data</b>	Radio tracking of select target species.
<b>Description of Methodology for Collecting Data</b>	Radio tracking using VHF or nanotags Best for short-range tracking (<25km).
<b>Methodology for Analyzing data</b>	<ul style="list-style-type: none"> <li>Depends on movement ecology of target species.</li> </ul>
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>If focused on birds from nearby breeding colony, throughout breeding season.</li> <li>Depends on biology of target species and battery life of transmitters.</li> </ul>
<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>Depends on biology of target species and range of transmitters and receiving stations.</li> </ul>
<b>Strengths</b>	<ul style="list-style-type: none"> <li>A network of receiving stations could potentially track movements of target species throughout a region.</li> <li>Could be used to track movements of nocturnal and diurnal migrants.</li> <li>Nanotags have the ability to track movements of small birds (e.g., passerines) and bats.</li> <li>Potential to accurately assess the position of individuals.</li> </ul>
<b>Limitations</b>	<ul style="list-style-type: none"> <li>Low sample size may not represent larger population.</li> <li>Absence of target individuals in the developed area does not necessarily mean that the population is not using the area.</li> <li>Trade-off between battery life and data collection.</li> <li>Data collection can be intensive with multiple observers (or boats) needed.</li> <li>Receiving stations general can detect transmitters within 10-20 km of</li> </ul>

	station.
<b>Cost</b>	<ul style="list-style-type: none"> <li>Moderate (depends on number of individuals tracked and their locations)</li> </ul>
<b>Type of Data Collected</b>	<ul style="list-style-type: none"> <li>Real time locations of target species</li> </ul>
<b>Data Output</b>	<ul style="list-style-type: none"> <li>Phenology of spatially-explicit movements of target species.</li> </ul>
<b>Examples where this technique has been used for ORED monitoring</b>	Perrow et al. 2006; Walls et al. 2009.

**Avian Protocol 9: Satellite Tracking.** Assess changes in avian movement ecology, such as migration and foraging flight paths between foraging and roosting sites due to ORE facility.

<b>Indicator(s) of Impact</b>	Changes in flight paths of foraging or commuting birds within development area.
<b>Methodology or Technique to Collect Data</b>	Satellite tracking.
<b>Description of Methodology or Technique(s) for Collecting Data</b>	<ul style="list-style-type: none"> <li>Position of individual is estimated on each satellite pass.</li> <li>Accurate to 250m, but can be variable in accuracy.</li> <li>Number of locations per day depends on programmed duty cycle and lat/longitude.</li> <li>Observers not required to track birds.</li> </ul>
<b>Methodology for Analyzing data</b>	<ul style="list-style-type: none"> <li>Home range analysis.</li> </ul>
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>Variable duty cycles can make transmitters last from 6 month to 2 years, depending on how often tag is turned on. Current recommendations suggest tags should be on for at least 4 hours to increase the probability of accurate fixes.</li> <li>Depends on biology of target species.</li> </ul>
<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>Development area</li> </ul>
<b>Strengths</b>	<ul style="list-style-type: none"> <li>Ideal for general studies of long-distance movements. Able to track movements over 1000s of kilometers.</li> </ul>

	<ul style="list-style-type: none"> <li>• Birds can be tracked in variable weather conditions.</li> <li>• Accurate location data of individuals.</li> <li>• Information on habitat preferences and larger scale movements.</li> </ul>
<b>Limitations</b>	<ul style="list-style-type: none"> <li>• More suited for large scale movements; not useful for fine-scale.</li> <li>• Depending on target species, transmitters may have to be surgically implanted; a veterinarian would need to be hired for this procedure.</li> <li>• Generally only about 700 fixes per battery cycle.</li> <li>• Low sample size may not represent larger population.</li> <li>• Absence of target individuals in the developed area does not necessarily mean that the population is not using the area.</li> <li>• Given current available transmitters, unsuitable for species that weight less than approx. 400 grams because current recommendations suggest that tags should not be more than 3% of body mass and the current smallest tags are about 12 g.</li> <li>• Trade-off between battery life and data collection.</li> </ul>
<b>Cost</b>	<ul style="list-style-type: none"> <li>• Low (Depends on number of individuals tracked)</li> </ul>
<b>Type of Data Collected</b>	<ul style="list-style-type: none"> <li>• Spatially-explicit location data, accurate to within 250 m of actual bird's location, with a time stamp accurate to the nearest second.</li> </ul>
<b>Data Output</b>	<ul style="list-style-type: none"> <li>• Spatially-explicit location data can be as accurate to within 250 m of actual bird's location.</li> </ul>
<b>Examples where this technique has been used for ORED monitoring</b>	Griffin et al. 2010; Walls et al. 2009.

**Avian Protocol 10: GPS Tracking.** Assess changes in avian movement ecology, such as migration and foraging flight paths between foraging and roosting sites due to ORE facility.

<b>Indicator(s) of Impact</b>	Changes in flight paths of foraging or commuting birds within development area.
<b>Methodology or Technique to Collect Data</b>	GPS tracking.
<b>Description of Methodology or Technique(s) for collecting data</b>	<ul style="list-style-type: none"> <li>• Accurate to within <math>\pm 5m</math></li> <li>• Some technologies require birds to be recaptured to upload data.</li> <li>• Works best on colony-breeding birds which can be easily recaptured.</li> </ul>
<b>Methodology for Analyzing data</b>	<ul style="list-style-type: none"> <li>• Before-After design, home range analysis.</li> </ul>
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>• Depends on biology of target species.</li> </ul>

<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>• Not relevant.</li> </ul>
<b>Strengths</b>	<ul style="list-style-type: none"> <li>• Extremely accurate location data of individuals compared other technologies.</li> <li>• Can track a bird for over one year, with thousands of fixes during this annual cycle.</li> <li>• Information on habitat preferences and movement.</li> </ul>
<b>Limitations</b>	<ul style="list-style-type: none"> <li>• Low sample size may not represent larger population.</li> <li>• Absence of target individuals in the developed area does not necessarily mean that the population is not using the area.</li> <li>• Mass of available tags makes this technology unsuitable for species less than 400 g, as current GPS technologies go down to 12 g.</li> <li>• Studies have shown that tags should not be more than 3% of body mass.</li> <li>• Trade-off between battery life and data collection.</li> </ul>
<b>Cost</b>	<ul style="list-style-type: none"> <li>• Low (Depends on number of individuals tracked)</li> </ul>
<b>Type of Data Collected</b>	<ul style="list-style-type: none"> <li>• Real time locations, accurate to &lt; 5 m to nearest minute.</li> </ul>
<b>Data output</b>	<ul style="list-style-type: none"> <li>• Spatially-explicit location data, can be as accurate to within 5 m of actual bird's location.</li> </ul>
<b>Examples where this technique has been used for ORED monitoring</b>	Walls et al. 2009.

**Avian Protocol 11: Thermal Animal Detection System.** Assess direct mortality (above-water collision) of marine birds due to ORE development.

<b>Indicator(s) of Impact</b>	Birds found dead or injured due to direct collision with infrastructure above the water.
<b>Methodology or Technique to Collect Data</b>	Thermal Animal Detection System (TADS)
<b>Description of Methodology or Technique(s) for collecting data</b>	<p>Thermal cameras use the heat radiating off of birds to create a thermal image.</p> <p>Operation should be limited to 1-2 km due to the low optical resolution of the thermal camera.</p>
<b>Monitoring Design and Analysis Recommendations</b>	<ul style="list-style-type: none"> <li>• Minimum of one TAD per wind facility with a maximum of 1 TAD per wind turbine.</li> </ul>

<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>• 24 hours per day/ 7 days per week for an entire year.</li> <li>• One to two years post construction.</li> </ul>
<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>• Individual wind turbines.</li> </ul>
<b>Strengths</b>	<ul style="list-style-type: none"> <li>• Ability to remotely monitor collision risk 24 hours per day, 7 days per week.</li> <li>• Can detect nocturnal targets including bats.</li> </ul>
<b>Limitations</b>	<ul style="list-style-type: none"> <li>• Low optical resolution makes identifying to species difficult.</li> <li>• Difficult in harsh offshore conditions.</li> <li>• Uncertainty of effectiveness in inclement weather.</li> </ul>
<b>Cost</b>	<ul style="list-style-type: none"> <li>• Low (depends on how many TADS are put in place)</li> </ul>
<b>Type of Data Collected</b>	<ul style="list-style-type: none"> <li>• Number of targets approaching and colliding with turbines.</li> </ul>
<b>Data Output</b>	<ul style="list-style-type: none"> <li>• Total number of targets and collisions near wind facility, including a time stamp.</li> </ul>
<b>Examples where this technique has been used for ORED monitoring</b>	<p>Walls et al. 2009; Desholm et al. 2004; Desholm et al.. 2006.</p>

**Avian Protocol 12: Sonar and Video Technology.** Assess direct mortality (under water-collision) of marine birds due to ORE development.

<b>Indicator(s) of Impact</b>	Birds found dead or injured due to direct collision with infrastructure under water.
<b>Methodology or Technique to Collect Data</b>	Point count could be used to Monitor avian use of project area from surface using a point count.  Currently, no remote technologies developed to detect underwater avian collisions (sonar and video technology developed for demersal fish strikes may work for birds).
<b>Description of Methodology or Technique(s) for Collecting Data</b>	Station observers to visually monitor the project area to determine if diving birds are using the project area.
<b>Monitoring Design and Analysis Recommendations</b>	<ul style="list-style-type: none"> <li>• Post-construction surveys.</li> </ul>
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>• Weekly during period when diving birds could be in study area.</li> <li>• One year post-construction.</li> </ul>
<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>• Project area.</li> </ul>
<b>Strengths</b>	<ul style="list-style-type: none"> <li>• Allow quick determination if potentially vulnerable species (diving birds) are using the study area.</li> <li>• Cost effective compared to other potential strategies</li> </ul>
<b>Limitations</b>	<ul style="list-style-type: none"> <li>• Practical only on days when observers could be stationed at project area</li> <li>• Only feasible for diurnal observations</li> <li>• Best for larger marine birds with high detection probabilities</li> </ul>
<b>Cost</b>	<ul style="list-style-type: none"> <li>• Moderate</li> </ul>
<b>Relationship to Other Protocols</b>	<ul style="list-style-type: none"> <li>• Can be conducted in combination with Fisheries Resources Protocol X4</li> </ul>
<b>Type of Data Collected</b>	<ul style="list-style-type: none"> <li>• Number of individuals of target species detected in study area.</li> </ul>
<b>Data Output</b>	<ul style="list-style-type: none"> <li>• Number of vulnerable targets (diving birds) in study area.</li> </ul>

<p><b>Examples where this technique has been used for ORED monitoring</b></p>	<p>None to our knowledge</p>
-------------------------------------------------------------------------------	------------------------------

**Marine mammals and sea turtles (MM&ST)**

**MM and/or ST Protocol 1: Visual surveys**

**Monitoring objectives:**

**Construction Noise (pile driving)/Decommissioning Noise (pile removal, explosives):** Mortality, injury, or disturbance of MM or ST by loud sounds

**Operational Noise:** Disturbance

**Vessel Collisions:** Mortality or injury

**Entanglements:** Mortality or injury

**Cable-laying:** Disturbance, mortality or injury

<p><b>Indicator(s) of the impact</b></p>	<p>Changes in local or regional distribution, abundance, or behavior of populations; Presence of dead or injured animals.</p>
<p><b>Summary of Methodology or Technique(s) to Collect Data</b></p>	<ul style="list-style-type: none"> <li>• Searching by trained observers for target species.</li> <li>• Typically observers are aboard ships and/or aircraft following pre-defined track-lines covering an area of interest, but surveys can be land-based for a specific focus (turtle nesting, pinniped rookeries or haul-outs).</li> <li>• Survey methodology can encompass higher-tech options, e.g., high-definition photography or videography.</li> <li>• Can secondarily provide data for mitigation, e.g., provide advance warning of animals nearby or approaching impact zone. (See Appendix B for more detailed descriptions)</li> </ul>
<p><b>Methodology for Analyzing data</b></p>	<ul style="list-style-type: none"> <li>• Line-transect or strip-transect analysis of survey data, using well-established methods, results in estimates of the density (and therefore abundance) of MM&amp;ST species within study area.</li> <li>• Since the range of each species is generally much larger than the scale of any given ORED project, the scope of these surveys will be too small to estimate population abundance.</li> <li>• There should also be BACI analysis on geospatial data given an appropriate sampling design.</li> </ul>

<b>Frequency and Duration</b>	Optimum would be at least 1 year prior to beginning construction, with the exact duration partially site-specific depending on the extent of prior sampling. Sampling should continue for the full duration of construction, and ideally at least 2-3 years post-construction. Survey frequency will be project-specific depending on the species present and their densities (rare species require more sampling to generate robust estimates).
<b>Spatial Scale</b>	The minimum scale would be the project area plus some buffer. For noise impacts, an acoustic propagation model will predict the maximum ranges of potential acoustic injury or disturbance; that will determine the minimum extent of the “impact” area for a survey. For effective BACI analysis, the “control” area should be beyond those ranges, but in an ecologically equivalent habitat with similar species and densities present.
<b>How well does this methodology account for environmental variability?</b>	Inter-annual variability in marine mammal and sea turtle populations is known to be high, and the duration of construction of any ORED facility will be much shorter than that variability. A well-designed BACI study with appropriate control and impact areas might be able to account for effects of variability in both habitat and population characteristics.
<b>Cost</b>	Moderate to High
<b>Other Considerations (E.g. Advantages or Disadvantages)</b>	Visual surveys only work well where population densities are sufficiently high to produce necessary sample sizes and statistically robust estimates; their use is also limited at night or under reduced visibility conditions (fog, high winds, storms).
<b>Relationship to other protocols</b>	Survey data can provide ground-truthing (confirmation of species IDs) for passive-acoustic monitoring (Protocol W2). Surveys can alert MMOs to animal presence (Protocol W3), or pass on observations of dead or injured animals to stranding responders for recovery and necropsy (Protocol W4).
<b>Data Format:</b>	There is a variety of existing formats for aerial or shipboard survey data, but they are effectively interchangeable if the necessary data fields are collected in the first place (Kenney, 2001, 2010; Halpin et al., 2009). The data collection and management methodology is sufficiently well-established so that any organization capable of fielding a survey effort is already familiar with data formats. Any additional standardization required can easily be established.

<b>Data Output:</b>	From the raw survey data, three basic types of data output are possible (see Kenney and Shoop, 2012, for a summary of aerial survey methods). At the most basic are geospatial data—sighting locations which can be mapped in GIS or summarized for statistical analysis. At the most rigorous level, estimates of species density can be computed using line-transect or strip-transect methodology, assuming that there are sufficient sightings of that species to generate the necessary sighting probability models. At the intermediate level of statistical rigor, it can be possible to develop relative abundance estimates (see Kenney and Vigness-Raposa, 2010 for an example).
<b>Examples where this methodology has been used:</b>	CETAP 1982; Waring et al., 2010; Allen and Angliss, 2011; Caretta et al., 2011; Forney, 2000; Ferguson et al., 2006a, 2006b; Redfern et al., 2006; Barlow et al., 2009; Becker et al., 2010; Teilmann et al., 2006a, 2006b; Thompson et al., 2010; Edrén et al., 2010; Malme et al., 1984; Frankel and Clark, 1998

**MM Protocol 2: Passive Acoustic Monitoring**

**Monitoring objectives:**

**Construction Noise (pile driving)/Decommissioning Noise (pile removal, explosives):** Disturbance of marine mammals by loud sounds

**Operational Noise:** Disturbance; Changes in distribution or abundance

**Vessel Traffic:** Disturbance; Changes in distribution or abundance

**Cable-laying:** Disturbance; Changes in distribution or abundance

<b>Indicator(s) of the impact</b>	Changes in distribution, abundance, or behavior of populations.
<b>Summary of Methodology to Collect Data</b>	Passive acoustic monitoring essentially involves listening to ambient sounds and identifying vocalizations produced by marine mammals.
<b>Methodology for Analyzing data</b>	BACI analysis on a variety of data metrics.
<b>Frequency and Duration</b>	Optimum would be at least 1 year prior to construction, with the exact duration partially site-specific depending on the extent of prior sampling. Sampling should continue for the full duration of construction, and ideally at least 2-3 years post-construction.

<b>Spatial Scale</b>	For noise impacts, an acoustic propagation model will predict the maximum ranges of potential acoustic injury or disturbance; that will determine the minimum extent of the “impact” area for a survey. For effective BACI analysis, the “control” area should be beyond those ranges, but in an ecologically equivalent habitat with similar species and densities present. Cost considerations will factor in to decisions on the number of sensors that can be deployed.
<b>How well does this methodology account for environmental variability?</b>	Inter-annual variability in marine mammal and sea turtle populations is known to be high, and the duration of construction of any ORED facility will be much shorter than that variability. A well-designed BACI study with appropriate control and impact areas might be able to account for effects of variability in both habitat and population characteristics.
<b>Cost</b>	Moderate to High
<b>Other Considerations (E.g. Advantages or Disadvantages)</b>	PAM only works on species that routinely vocalize (i.e., not on seals or sea turtles). Autonomous sensors that must be recovered to download the data provide no real-time monitoring capability. Because of limitations on data uplink bandwidth, typical near-real-time sensors provide only detections of pre-programmed species (usually right whales) and not multi-species data or continuous data.
<b>Relationship to other protocols</b>	Data can feed into other MM&ST protocols.
<b>Data Format</b>	The data output from passive-acoustic monitoring will depend heavily on the sensors and sampling methodology employed
<b>Data Output</b>	<p>Continuous data can be analyzed for all species that might be present and whose vocalizations fall within the frequency range recorded.</p> <p>Actual tracks of vocalizing individuals can be compared between control and impact areas or times.</p> <p>Combined visual data from shipboard surveys and simultaneous PAM data from towed arrays can be used to derive density estimates</p> <p>Porpoise-positive minutes (minutes with clicks recorded), waiting time between encounters (detections of sets of clicks), waiting time from the end of pile-driving to the first detection, duration of encounters, and number of clicks per porpoise-positive minute (e.g. Carstensen et al., 2006; Teilmann et al., 2006a; Tougaard et al., 2009a)</p>

<b>Examples where this methodology has been used:</b>	Carstensen et al., 2006; Teilmann et al., 2006a, 2008; Diederichs et al., 2008; Tougaard et al., 2009a; Clausen et al., 2010; Brandt et al., 2011; Tyack et al., 2011; NMFS, 2010d, 2010e; Risch et al., 2012
-------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

**MM&ST Protocol 3: Marine Mammal Observers**

**Monitoring objectives:**

**Construction Noise (pile driving)/Decommissioning Noise (pile removal, explosives):** Mortality, injury, or disturbance of marine mammals or sea turtles

**Vessel Traffic:** Mortality, injury, or disturbance of marine mammals or sea turtles

**Cable-laying:** Mortality, injury, or disturbance of marine mammals or sea turtles

**Entanglement:** Mortality or injury of marine mammals or sea turtles

<b>Indicator(s) of the impact</b>	Presence of dead or injured animals; detection of animals within impact zones of potentially harmful activities.
<b>Summary of Methodology or Technique(s) to Collect Data</b>	Marine Mammal (or Protected Species) Observers are trained observers posted on board vessels in an active construction or operational area. The primary objective of an MMO program often is mitigation—detection of animals in potential zones of injury and shutting down operation and/or stopping or diverting vessels.
<b>Methodology for Analyzing data</b>	List of animals observed within given ranges of the activity being monitored and any observed behavioral reactions.
<b>Frequency and Duration</b>	MMOs should be deployed continuously for the full duration of construction, as well as on board vessels where the risk of impacts is high, which is project-specific.
<b>Spatial Scale</b>	Limited to the visual range of an observer.
<b>Cost</b>	Low
<b>Other Considerations (E.g. Advantages or Disadvantages)</b>	Effectiveness is limited by visibility and distance. For example noise disturbance from pile-driving is possible beyond the distance where an MMO posted near the construction site might see an animal.
<b>Relationship to other protocols</b>	Data from visual and real-time passive acoustic surveys (Protocols W1 and W2) can alert MMOs to animals near or approaching the impact zone.

<b>Data Format</b>	Standard sighting data (date, time, location, species, numbers, behaviors), which can be added to datasets from any survey programs  Behavioral observations can also be used to assess potential negative effects of project activities on behavior.
<b>Examples where this methodology has been used:</b>	MMS, 2009; NMFS, 2003; NMFS, 2010c; DON, 2009b

**MM&ST Protocol 4: Stranding Response Networks**

**Monitoring objectives:**

**Construction Noise (pile driving)/Decommissioning Noise (pile removal, explosives):** Mortality or injury of marine mammals or sea turtles

**Vessel Traffic:** Mortality or injury of marine mammals or sea turtles

**Cable-laying:** Mortality or injury of marine mammals or sea turtles

**Entanglement:** Mortality or injury of marine mammals or sea turtles

<b>Indicator(s) of the impact</b>	Detection of dead or injured animals with evidence of causation from impacts of the project.
<b>Summary of Methodology to Collect Data</b>	Visual survey; Passive acoustic monitoring; MMOs  Stranding network; Tagging; Stress hormones
<b>Methodology for Analyzing data</b>	Standard veterinary pathology methods.
<b>Frequency and Duration</b>	Continuous for the duration of the project.
<b>Spatial Scale</b>	Coast-wide with enhanced response in regions where ORED facilities are planned, under construction, or in operation.
<b>Cost</b>	Low
<b>Other Considerations (E.g. Advantages or Disadvantages)</b>	Stranding networks are already in place and would require only enhancement to be effective for monitoring purposes.
<b>Relationship to other</b>	Data feed into other MM&ST protocols.

<b>protocols</b>	
<b>Data Output:</b>	Cause-of-death determinations for all marine mammals or sea turtles found dead in the vicinity of ORED facilities, based on standard veterinary necropsy and pathology methods
<b>Examples where this methodology has been used:</b>	Waring et al., 2010; Allen and Angliss, 2011; Caretta et al., 2011

**MM&ST Protocol 5: Tagging**

**Monitoring objective:** Disturbance of marine mammals or sea turtles by noise, activities, or structures from ORE development.

<b>Indicator(s) of the impact</b>	Detection of changes in fine-scale distribution, movement, or behavior of individuals.
<b>Summary of Methodology or Technique(s) to Collect Data</b>	Tagging involves fixing a device to an individual animal and then tracking the location of that individual, often recording other data parameters simultaneously.
<b>Methodology for Analyzing data</b>	Highly dependent on the type of tag used. BACI or impact gradient analysis on geospatial data and/or behavioral data can be possible—assuming sufficient sample sizes and that the tagged animals cooperate by utilizing an appropriate selection of habitat sites.
<b>Frequency and Duration</b>	Species- and project-specific; a power analysis would help to define the number and duration of tag deployments necessary to produce statistically reliable results.
<b>Spatial Scale</b>	Species- and project-specific; each tagged animal will actually define its own spatial scale. Given the likelihood of an individual tagged animal moving far beyond the boundaries of any given project study area; a large-scale tagging/telemetry study may be one of the better methods for addressing cumulative impacts of multiple MRE projects along broad areas of the U.S. coastline.
<b>Cost</b>	Moderate to High
<b>Other Considerations (E.g. Advantages or Disadvantages)</b>	Tagging is logistically challenging, entails high costs, and poses some risk to the animals; in addition it can be difficult to generate effective sample sizes, with the expectation that some proportion of tagged individuals will leave the study area.

<b>Relationship to other protocols</b>	Tagging is similar to the stress hormone protocol (Protocol 6) in that it gets more into effects research that might be conducted if other monitoring results suggest that there might be effects.
<b>Data Format:</b>	The type of data resulting from tagging studies can be extremely variable, and will be dependent upon the type of tags employed.
<b>Data Output:</b>	All tagging will result in some level of geospatial data—locations of the tagged individual at particular times. Depending on the tag, these can range from simply deployment and recovery locations for flipper tags to small numbers of locations per day for satellite or geo-locator tags, to detailed movement tracks for GPS archival tags. Telemetry tags with depth sensors to monitor diving behavior can provide simple data summaries (e.g., number of dives in the previous 24 hr, maximum depth) for tags with restricted reporting bandwidth to detailed, continuous dive profiles for days to months in the case of archival tags. Methods have been developed for taking the depth and accelerometer data from DTAGs and deriving 3-dimensional graphics or even animations of the submerged foraging behavior of tagged whales (Ware et al., 2006).
<b>Examples where methodology has been used:</b>	Tougard et al., 2003; Teilmann et al., 2006b; Müller and Adelung, 2008; Friedlander et al. 2009; Miller et al. 2009; Nowacek et al, 2004

**MM & ST Protocol 6: Stress Hormone Assessment**

**Monitoring objective:** Quantification of physiological stress related to disturbance from ORE development activities.

<b>Indicator(s) of the impact</b>	Elevated levels of stress-related corticosteroid hormones in animals subject to disturbance from activities associated with ORED development.
<b>Summary of Methodology or Technique(s) to Collect Data</b>	Measurement of levels of corticosteroid hormones and/or their metabolites is a standard biomedical technique. Data collection would be by collecting fecal or blow samples from free-swimming whales. Sampling could be in a control-impact design or a gradient design along a continuum of distances from a potential disturbance.
<b>Methodology for Analyzing data</b>	Standard hormone bio-assays; statistical comparison of levels between control and treatment groups.
<b>Frequency and Duration</b>	To be defined by the number of samples necessary to obtain statistically meaningful results.
<b>Spatial Scale</b>	Project-specific, depending on the ranges at which potential disturbance has been detected by other monitoring studies.
<b>Cost</b>	Moderate
<b>Other Considerations (E.g. Advantages or Disadvantages)</b>	Stress hormone assessment on free-swimming marine mammals is a relatively new method that to date has only been applied to a couple of large whale species, however those are usually the species of the greatest conservation concern.
<b>Relationship to other protocols</b>	This methodology is more focused research that would not likely be employed until other monitoring (e.g., surveys, PAM) has detected changes in distribution related to the project. Also see the tagging protocol (Protocol 5). A sort of controlled exposure experiment could be conducted by sampling from animals tagged with DTAGs or similar tags that monitor received levels of sound, and correlating stress hormone levels with noise exposures.
<b>Data Output:</b>	Stress hormone concentrations from samples collected in control and impact areas or from along an exposure gradient.
<b>Examples where methodology has been used:</b>	Rolland et al. 2007; Hogg et al. 2009; Rolland et al. 2012

**MM&ST Protocol 7: Underwater Photography**

**Monitoring objective:** Disturbance or loss of habitat of sea turtles by device removal during decommissioning of ORE devices.

<b>Indicator(s) of the impact</b>	Disturbance of animals during cable or device removal; detection of dead or injured animals
<b>Summary of Methodology to Collect Data</b>	Underwater camera mounted on ORED structure(s) to collect time-lapse photography of “resident” sea turtles and marine mammals prior to structure(s) decommissioning
<b>Methodology for Analyzing data</b>	Direct reporting of data; qualitative analysis to detect presence/absence of marine mammals or turtles
<b>Frequency and Duration</b>	Begin monitoring 1 month prior to cable-laying/removal or decommissioning of structure(s), on day of laying/removal, and 1 month following removal; avoid nighttime removals. Could be combined with fisheries ROV/SCUBA survey efforts
<b>Spatial Scale</b>	Small: Area immediately surrounding structure to be removed
<b>Cost</b>	Low
<b>Other Considerations (E.g. Advantages or Disadvantages)</b>	Time-lapse underwater photography is preferred method, since it is unobtrusive and provides more complete coverage.
<b>Relationship to other protocols</b>	Can be combined with Fisheries Protocol 3
<b>Data Format:</b>	Individual sightings records, including species identification, size estimates (if possible), and behavioral characteristics noted.
<b>Data Output:</b>	Video record and direct reporting of individual “resident” marine mammals and turtles in vicinity of ORED structures would trigger mitigation actions to disperse/relocate/fire warning charges to prevent impacts.
<b>Examples where methodology has been used:</b>	Rosman et al., 1987; Klima et al., 1988

**MM&ST Protocol 8: SCUBA Surveys**

**Monitoring objective:** Disturbance of sea turtles during cable installation or removal; Disturbance or loss of habitat of sea turtles by ORE device removal during decommissioning.

<b>Indicator(s) of the impact</b>	Disturbance of sea turtles during cable installation/removal; Disturbance of animals during cable or device removal; detection of dead or injured animals
<b>Summary of Methodology to Collect Data</b>	SCUBA diver surveys (e.g. during cable laying/removal or around structures to be decommissioned/removed)
<b>Methodology for Analyzing data</b>	Direct reporting of data; qualitative analysis to detect presence/absence of marine mammals or turtles
<b>Frequency and Duration</b>	Begin monitoring 1 month prior to cable-laying/removal or decommissioning of structure(s), on day of laying/removal, and 1 month following removal; avoid nighttime removals. Could be combined with fisheries ROV/SCUBA survey efforts
<b>Spatial Scale</b>	Small: Area immediately surrounding structure to be removed
<b>Cost</b>	Low
<b>Other Considerations (E.g. Advantages or Disadvantages)</b>	Diver surveys are useful, but can miss turtles, likely due to submersible/observer presence
<b>Relationship to other protocols</b>	Can be combined with Marine Mammal and Sea Turtle Protocol 7 and 9; Fisheries Protocol 3
<b>Data Format:</b>	Individual sightings records, including species identification, size estimates (if possible), and behavioral characteristics noted.
<b>Data Output:</b>	Direct reporting of individual marine mammals and turtles in vicinity of cable-laying or resident animals at ORED structures would trigger mitigation actions to disperse/relocate/fire warning charges to prevent impacts.
<b>Examples where this methodology has been used:</b>	Klima et al., 1988; Rosman et al. 1987

**MM&ST Protocol 9: ROV Surveys**

**Monitoring objective:** Disturbance of sea turtles during cable installation or removal; Disturbance or loss of habitat of sea turtles by ORE device removal during decommissioning.

<b>Indicator(s) of the impact</b>	Disturbance of sea turtles during cable installation/removal; Disturbance of animals during cable or device removal; detection of dead or injured animals
<b>Summary of Methodology or Technique(s) to Collect Data</b>	Video surveys with ROV (e.g. during cable laying/removal or around structures to be decommissioned/removed)  Survey areas directly surrounding structures to be removed or along the path where cable-laying/removal will occur
<b>Methodology for Analyzing data</b>	Direct reporting of data; qualitative analysis to detect presence/absence of marine mammals or turtles
<b>Frequency and Duration</b>	Begin monitoring 1 month prior to cable-laying/removal or decommissioning of structure(s), on day of laying/removal, and 1 month following removal; avoid nighttime removals. Could be combined with fisheries ROV survey efforts
<b>Spatial Scale</b>	Small: Area immediately surrounding structure to be removed or ahead of jet plow along path of cable laying/removal
<b>Cost</b>	Low if combined with fisheries surveys; moderate if conducted independently
<b>Other Considerations (E.g. Advantages or Disadvantages)</b>	ROV surveys can miss animals, likely due to submersible/observer presence or underwater visibility considerations, but are not weather/sea state-dependent.
<b>Relationship to other protocols:</b>	Can be combined with Marine Mammal and Sea Turtle Protocol W7 and W8; Fisheries Protocol W3
<b>Data Format:</b>	Individual sightings records, including species identification, size estimates (if possible), and behavioral characteristics noted.
<b>Data Output:</b>	Direct reporting of individual marine mammals and turtles in vicinity of cable-laying or resident animals at MRE structures would trigger mitigation actions to disperse/relocate/fire warning charges to prevent impacts.
<b>Examples where this methodology has been</b>	Rosman et al. 1987

<b>used:</b>	
--------------	--

**Fish**

**Fisheries Protocol 1a- Trawl Surveys.** Monitor for changes in meso-scale distribution and abundance of fish species in the vicinity of an ORE installation.

<b>Indicator(s) of the impact</b>	<ul style="list-style-type: none"> <li>• Shift in fish distribution or abundance overall or on a seasonal basis</li> <li>• Shift in species composition</li> <li>• Increase or decrease in catchability (Catch per Unit Effort) of commercially or recreationally targeted species</li> </ul>
<b>Methodology or Technique to Collect Data</b>	Otter Trawl Survey
<b>Description of Methodology or Technique(s) for collecting data</b>	<ul style="list-style-type: none"> <li>• Using BACI design with multiple control locations no less than 1km outside of the project area</li> <li>• Control locations selected to have similar bottom types and benthic habitat as project area trawl locations</li> <li>• Trawl locations from random station grid</li> <li>• Surveys conducted a minimum of four times/year</li> <li>• Baseline trawl locations and paths will be selected to be able to follow the same route after construction</li> <li>• All fish species sampled, with particular attention paid to commercially, recreationally, and ecologically important species</li> <li>• Sampling of weight and length of species</li> <li>• One inch knotless cod end liner</li> <li>• Trawl speed of 3 knots</li> <li>• Trawl duration 20 minutes</li> </ul>
<b>Methodology for Analyzing data</b>	ANOVA on numbers of individuals, size and weight distribution; multivariate analysis of catch/community composition, multidimensional scaling, cluster analysis (Primer-E)
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>• 2 years of baseline data in pre-construction period (surveys at least 4 times/year both years)</li> <li>• 4 surveys/year during post-construction for minimum of 5 years</li> </ul>
<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>• Random stratified surveys selected from the following stratification: 10 sites within .5 km of renewable energy site; 10 sites between .5-2.5 km of renewable energy site; 10 control sites (at greater than 2.5 km from site)</li> <li>• Control sites should be selected from areas with similar bathymetry and bottom type to renewable energy site</li> <li>• A minimum of 30 trawls per survey period</li> </ul>

<b>How well does this methodology account for environmental variability?</b>	Seasonal and interannual variability
<b>Cost</b>	Moderate (depends on number of surveys)
<b>Other Considerations (E.g. Advantages or Disadvantages)</b>	<ul style="list-style-type: none"> <li>• Not all survey types and gear types will be appropriate to each location. The gear and survey types should be selected based on the issues of greatest concern.</li> <li>• Trawl survey will sample mostly demersal species rather than pelagic species. Survey limited to those species most prone to be caught in the net, and will under-sample some species, e.g. lobsters and crabs.</li> <li>• The commercial fishing industry should be consulted on the type of gear used.</li> <li>• The commercial fishing industry should be involved in data collection and survey design when feasible, including the selection of trawl stations.</li> </ul>
<b>Relationship to Other Protocols:</b>	Can be combined with Fisheries Protocol X1b or X2
<b>Data Format:</b>	<ul style="list-style-type: none"> <li>• Total individuals/area</li> <li>• Total biomass/area</li> <li>• # Individuals per species and area</li> <li>• Biomass per species and area</li> <li>• Diversity</li> <li>• Length frequency distribution of dominant species</li> <li>• Time series</li> </ul>
<b>Data Output:</b>	Catch per unit effort (CPUE) for total catch and on a species level; community dynamics
<b>Examples where this methodology has been used:</b>	Bundesamt für Seeschifffahrt und Hydrographie, 2007; Bonzek et al. 2008; CEFAS 2004.

**Fisheries Protocol 1b - Ventless Trap Surveys.** Monitor for changes in distribution and abundance of lobster/crab species or some fish species in the vicinity of an ORE installation.

<b>Indicator(s) of the impact</b>	<ul style="list-style-type: none"> <li>• Shift in distribution (of lobster, crab, rock fish) or abundance overall or on a seasonal basis</li> <li>• Shift in species composition</li> <li>• Increase or decrease in catchability (Catch per Unit Effort) of species by commercial fishing gear</li> </ul>
-----------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

<b>Methodology or Technique to Collect Data</b>	Fixed Gear Survey with Ventless Traps
<b>Description of Methodology or Technique(s) for collecting data</b>	<ul style="list-style-type: none"> <li>• Using BACI design with multiple control locations</li> <li>• Surveys conducted in spring and fall</li> <li>• Control locations selected to have similar bottom types and benthic habitat as project area trawl locations</li> <li>• Sampling of weight and length of species</li> </ul>
<b>Methodology for Analyzing data</b>	ANOVA on numbers of individuals, size and weight distribution; multivariate analysis of catch/community composition
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>• 2 years of baseline data in pre-construction period (seasonal surveys 4 x/year both years)</li> <li>• Seasonal (4/year) during post-construction for minimum of 5 years</li> </ul>
<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>• Traps set within the renewable energy installation, and at random stratified sites at varying distances from the renewable energy site (e.g. 1 km, 10 km, 25 km)</li> </ul>
<b>How well does this methodology account for environmental variability?</b>	Seasonal and interannual variability
<b>Cost</b>	Low
<b>Other Considerations (E.g. Advantages or Disadvantages)</b>	<ul style="list-style-type: none"> <li>• Not all survey types and gear types will be appropriate to each location. The gear and survey types should be selected based on the issues of greatest concern.</li> <li>• The gear and techniques used by the commercial fishing industry should be mirrored in the survey design when sampling commercially-important fish species.</li> <li>• The commercial fishing industry should be involved in data collection and survey design when feasible.</li> <li>• While ventless trap surveys are often used for crustaceans, they may be useful for species such as black sea bass, rock fish, or other species that are attracted to structures and can be caught by traps or pots.</li> </ul>
<b>Relationship to Other Protocols:</b>	Can be combined with Fisheries Protocol X1a or X2
<b>Data Format:</b>	<ul style="list-style-type: none"> <li>• Total individuals/area</li> <li>• Total biomass/area</li> <li>• # Individuals per species and area</li> <li>• Biomass per species and area</li> <li>• Length frequency distribution of dominant species</li> </ul>

<b>Data Output:</b>	Catch per Unit Effort (CPUE) at species level
<b>Examples where methodology has been used:</b>	Maine Department of Marine Fisheries. 2006. Courchene, B., and K.D.E. Stokesbury. 2011.

**Fisheries Protocol 2- Monitoring for project-scale changes**

**Monitoring objectives:** Monitor for micro-scale changes in abundance or species composition of fish around ORE structures or along cable routes, including non-native species, resulting from disturbance (from noise, presence of devices), or attraction to ORE devices (aggregation or reef effects). Also, monitor for changes in catchability of commercially and recreationally targeted fish in the vicinity of the ORE installation.

<b>Indicator(s) of the impact</b>	<ul style="list-style-type: none"> <li>- Increase or decrease in fish abundance</li> <li>- Increase or decrease in target species</li> <li>- Shift in species composition</li> <li>- Increase in presence of non-native species</li> <li>- Increase or decrease in catchability (Catch per Unit Effort) of commercially or recreationally targeted species</li> </ul>
<b>Methodology or Technique to Collect Data</b>	Gillnet surveys and/or trammel net surveys and/or beam trawl surveys
<b>Description of Methodology or Technique(s) for collecting data</b>	<p><u>Gillnet or trammel net surveys:</u></p> <ul style="list-style-type: none"> <li>• Installation-based surveys a minimum of 6 days/year</li> <li>• Three deployments each spring and fall for 1-2 days each</li> <li>• Installation at a minimum of three locations within footprint of renewable energy facility, and three reference locations in similar habitat, no less than 1km from footprint sites</li> </ul> <p><u>Beam trawl surveys:</u></p> <ul style="list-style-type: none"> <li>• Seasonal tows (spring, summer, fall, winter)</li> <li>• minimum of 3 locations within the footprint of the installation (between devices) - if possible</li> <li>• 9 ft. beam trawl with 1 in. knotless liner recommended</li> <li>• Tows at a minimum of three locations within footprint of renewable energy facility, and three reference locations in similar habitat, no less than 1km from footprint sites</li> </ul>

	Survey area can be expanded to include cable route, particularly when electromagneto-sensitive species (e.g. elasmobranchs) are of concern
<b>Methodology for Analyzing data</b>	ANOVA on # species, # of fish, multivariate analysis of fish community characteristics (Primer-E), multidimensional scaling, cluster analysis
<b>Frequency and Duration</b>	Baseline survey pre-construction (4 surveys, one each in spring, summer, fall, and winter)  Seasonal (4 times/year) during operation for 3 years
<b>Spatial Scale</b>	<u>Gillnets/Trammel nets</u> : Minimum of three installations within renewable energy footprint, and an equal number of reference stations in similar habitat  <u>Beam trawl</u> : Minimum of three locations within renewable energy footprint, and an equal number of reference stations in similar habitat
<b>How well does this methodology account for environmental variability?</b>	Seasonal and interannual variability
<b>Cost</b>	Low

<p><b>Other Considerations (E.g. Advantages or Disadvantages)</b></p>	<p>Gear type(s) used for the survey should depend on the fish species under consideration (commercially/recreationally important species, species of conservation importance), and the gear type that will be most effective in assessing changes to the abundance and distribution of these species on a fine scale.</p> <p>Gillnet surveys will undersample demersal species but can sample pelagic species, which are difficult to sample by other means.</p> <p>Gillnets are fairly size selective and will not provide a good estimate of overall biomass of the area. Combining gillnet and beam trawl surveys can account for a larger spectrum of fish species.</p> <p>Trammel nets can capture more fish than gillnets and will provide a greater picture of size distribution. However, trammel nets can be highly destructive and need to be checked or removed frequently.</p> <p>Passive nets can be deployed much closer to the devices than active trawling.</p> <p>Beam trawls can supplement otter trawls by trawling within an offshore renewable energy installation or between devices to sample within the footprint of a project, where otter trawling may not be feasible</p> <p>Beam trawls can also sample harder bottom habitats and are more effective at assessing benthic invertebrates (e.g. scallops, lobsters, clams, crabs)</p>
<p><b>Relationship to Other Protocols:</b></p>	<p>Can be combined with Fisheries Protocol X1a or X1b</p>
<p><b>Data Format:</b></p>	<p><u>Gillnet/trammel net:</u></p> <p>Catch per unit effort (CPUE)</p> <p># Individuals per species and area</p> <p>Diversity</p> <p>Length frequency distribution of dominant and/or vulnerable species</p> <p><u>Beam trawl:</u></p> <p>Total individuals/area</p> <p>Total biomass/area</p>

	<p># Individuals per species and area</p> <p>Biomass per species and area</p> <p>Diversity</p> <p>Length frequency distribution of dominant and/or vulnerable species</p>
<b>Data Output:</b>	time series values for # of individuals, biomass, fish community composition, and species-specific length frequency; presence absence of non-native species
<b>Examples where this methodology has been used:</b>	Bundesamt für Seeschifffahrt Hydrographie 2007; CEFAS 2004

**Fisheries Protocol 3a and 3b- Reef and Aggregation Effects.** Monitor for changes in abundance or species composition of fish around ORE structures, including non-native species.

	<b>3a: Depth of installation &lt; 20 m</b>	<b>3b: Depth of installation &gt; 20 m</b>
<b>Indicator(s) of the impact</b>	<ul style="list-style-type: none"> <li>• Increase in fish abundance overall or in some species</li> <li>• Shift in species composition</li> <li>• Increase in presence of non-native species</li> </ul>	
<b>Methodology to Collect Data</b>	Video surveys with ROV	Visual surveys with SCUBA
<b>Description of Methodology or Technique(s) for collecting data</b>	<ul style="list-style-type: none"> <li>• Minimum of four devices, four transects/device.</li> <li>• Transects of 1 km, radiating out from devices in four directions</li> <li>• Lasers for measuring length of fish species</li> </ul>	<ul style="list-style-type: none"> <li>• Minimum of four devices, four transects/device.</li> <li>• Transects 1-5 m and 20 m from devices on four sides</li> <li>• Transects radiating out from devices in four directions</li> <li>• Transects of 15-30 min duration</li> <li>• Estimation of species length</li> </ul>

<b>Methodology for Analyzing data</b>	ANOVA on # species, # of fish, multivariate analysis of fish community characteristics (Primer-E), multidimensional scaling, cluster analysis	ANOVA on # species, # of fish, multivariate analysis of fish community characteristics (Primer-E), multidimensional scaling, cluster analysis
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>• 4 seasonal baseline surveys during pre-construction</li> <li>• seasonal surveys 4x/yr for 5 yrs minimum during operation</li> </ul>	<ul style="list-style-type: none"> <li>• 2 baseline surveys during pre-construction (spring and fall)</li> <li>• seasonal surveys 2x/yr for 5 yrs minimum during operation</li> </ul>
<b>Spatial Scale</b>	Small to medium	Small
<b>How well does this methodology account for environmental variability?</b>	Seasonal and interannual variability	Seasonal and interannual variability
<b>Cost</b>	Low (depends # of surveys)	Low
<b>Other Considerations (E.g. Advantages or Disadvantages)</b>	Can be combined with reef effect protocol for benthic habitat	Can be combined with reef effect protocol for benthic habitat
<b>Relationship to Other Protocols:</b>	Can be combined with Fisheries Protocol 2 or Marine Mammal and Sea Turtle Protocols 6-8 or Benthic Environment Monitoring Protocol 2	
<b>Data Format:</b>	<p># Species per area</p> <p># Individuals per species and area</p> <p>Change in species/individuals with distance from devices</p> <p>Biomass per species and area</p> <p>Length frequency distribution of dominant and/or vulnerable species</p>	
<b>Data Output:</b>	time series values for # of species, # individuals; biomass estimates; presence absence of non-native species	
<b>Examples where this methodology has been used:</b>	Rademacher, K.R. and J.H. Render. 2003; Love, M, J. Hyland, A. Ebeling, T. Herrlinger, A. Brooks, E. Imamura. 1994.	Wilhelmsson, D., T. Malm, and M. Ohman. 2006.

**Fisheries Protocol 4: Blade Strikes.** Monitor for blade strikes from tidal energy devices.

<b>Indicator(s) of the impact</b>	Observation of blade strike incidents
<b>Methodology to Collect Data</b>	Video or sonar surveys of tidal turbine
<b>Description of Methodology or Technique(s) for collecting data</b>	<ul style="list-style-type: none"> <li>• Video cameras or DIDSON sonar system installed on a subset of turbines (3-5)</li> <li>• Video or Sonar will detect the movement of fish in the immediate vicinity of the tidal turbine</li> <li>• Video or Sonar will detect the number of fish approaching the turbine and the number of fish that pass through the blades, both while the turbines are operating and at periods of slack tide</li> </ul>
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>• Sonar installation will occur twice during the first year (spring and fall) for X days at a time</li> </ul>
<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>• Subset of turbines (3-5) dispersed throughout turbine field</li> </ul>
<b>How well does this methodology account for environmental variability?</b>	Some seasonal variability
<b>Cost</b>	Moderate
<b>Other Considerations (E.g. Advantages or Disadvantages)</b>	<ul style="list-style-type: none"> <li>• One study (Verdant Power 2010) found the DIDSON system could not be continuously deployed because of biofouling and siltation. The system should be deployed for a set period of time and then removed.</li> <li>• The sonar system may not be useful for identifying fish at the species level.</li> <li>• Verdant Power (2010) found the DIDSON system useful where the water was too turbid for video monitoring; video monitoring may be more practical where turbidity is less</li> <li>• DIDSON system was only effective at a distance of 15 m for appropriate resolution</li> </ul>
<b>Relationship to Other Protocols:</b>	Can be combined with Avian Species Monitoring Protocol 12 or Fisheries Protocol 2

<b>Data Format:</b>	<ul style="list-style-type: none"> <li>• Frequency of targets per time (fish within 10m, 5 m, 1 m of turbine; fish passing through turbine)</li> <li>• Distribution of fish in vicinity of turbine in various environmental conditions (tidal movement, slack tide, day/night)</li> <li>• Presence/absence of fish in the vicinity of the device</li> <li>• Number of observed blade strikes or fish passing through the devices per unit of time (at different times of day and in different seasons)</li> </ul>
<b>Data Output:</b>	<p>Video stills (extracted from video footage)</p> <p>Sonar stills (extracted from sonar footage)</p>
<b>Examples where methodology has been used:</b>	Verdant Power 2010.

**Fisheries Protocol 5 – Spatial Use of Fishing Activity.** Monitor for changes in the spatial distribution of fishing activity (commercial and recreational) around ORE installation.

Protocol	5a	5b
<b>Indicator(s) of the impact</b>	<ul style="list-style-type: none"> <li>• Changes in numbers of vessels fishing near or inside of the ORE area (more or fewer vessels)</li> <li>• Change in the presence of fixed fishing gear (gillnets, pots, traps) inside of or around ORE installation (more or less fixed gear)</li> </ul>	
<b>Methodology or Technique to Collect Data</b>	VMS installed on vessels to track movements	Vessel surveys to count numbers of vessels fishing, fixed fishing gear in use
<b>Description of Methodology or Technique(s) for collecting data</b>	<ul style="list-style-type: none"> <li>• VMS systems installed on a sizeable and representative subsample of fishing fleet (e.g. 25% of vessels) known to utilize area where renewable energy infrastructure being installed</li> <li>• Analysis of VMS data from NMFS on vessels already installed with the device for the same time period</li> <li>• Movements of vessels tracked for 2 years pre-construction, during construction, and minimum 5 years post-construction</li> <li>• Tracking of movements with</li> </ul>	<ul style="list-style-type: none"> <li>• Transects with a boat to count numbers of fishing boats engaged in fishing inside and outside of renewable energy installation, including type of vessel, relative size of vessel, and type of fishing activity (type of gear; steaming, trawling, setting gear, etc.)</li> <li>• Transects with a boat to count numbers of fixed fishing gear, including gillnets, lobster traps, fish pots, etc.</li> <li>• Equal transects in a control area of equal size</li> </ul>

	VMS should also take place in a control area with no development, to exclude effects of fish movements, environmental variables, etc.	
<b>Methodology for Analyzing data</b>	GIS, Multidimensional scaling	GIS, Multidimensional scaling
<b>Frequency and Duration</b>	<ul style="list-style-type: none"> <li>• Year-round survey of vessel movement</li> <li>• 2 years pre-construction, during construction, and 5 years post-construction</li> </ul>	<ul style="list-style-type: none"> <li>• Year-round survey of fixed fishing gear and vessel activity</li> <li>• 2 years pre-construction, during construction, and 5 years post-construction</li> </ul>
<b>Spatial Scale</b>	<ul style="list-style-type: none"> <li>• Encompass entire renewable energy zone and large buffer area around renewable energy installation</li> <li>• Equal transects in a control area of equal size</li> </ul>	<ul style="list-style-type: none"> <li>• Encompass entire renewable energy zone and large buffer area around renewable energy installation</li> <li>• Equal transects in a control area of equal size</li> </ul>
<b>How well does this methodology account for environmental variability?</b>	Accounts for seasonal variability, and somewhat for interannual variability	Accounts for seasonal variability, and somewhat for interannual variability
<b>Cost</b>	Moderate (depends on the number of VMS units required)	Low
<b>Other Considerations (E.g. Advantages or Disadvantages)</b>	<ul style="list-style-type: none"> <li>• Some vessels will already be installed with VMS and are reporting VMS to NMFS. Only certain fisheries or vessels over a certain size are required to carry VMS systems</li> <li>• VMS should be installed on a variety of types of fishing boats engaged in a wide variety of fisheries to analyze which fisheries are most affected by the renewable energy installation</li> <li>• Analysis should be combined with analysis of trawl and fixed gear surveys from within and outside of renewable energy field.</li> </ul>	<ul style="list-style-type: none"> <li>• This methodology will be less expensive than installing VMS systems (depending on the number of vessels/VMS systems to be installed).</li> <li>• The most appropriate methodology may depend on the important fisheries within the area. When</li> <li>• Mobile fishing gear is predominant, VMS may be more suitable. When fixed fishing gear is predominant, transect surveys will be sufficient</li> <li>• Transects may be able to be combined with those for marine mammals or birds</li> <li>• Analysis should be combined</li> </ul>

		with analysis of trawl and fixed gear surveys from within and outside of renewable energy field.
<b>Relationship to Other Protocols:</b>	Can also be combined with Avian Species Monitoring Protocol 1 and Marine Mammal and Sea Turtles Monitoring Protocol 1	
<b>Data Format:</b>	Vessels/area Distribution of vessel types	
<b>Data Output:</b>	Spatial Use Maps	
<b>Examples where methodology has been used:</b>	Wiley, D.N.; Moller, J.C.; Zilinskas, K.A.. 2003.	

**Cultural and Historical Resources Task 1 – Geophysical surveys.**

For geophysical surveying, a two-tier approach is outlined. Tier 1 describes surveys, instrumentation, techniques, and resolution needed to achieve broad baseline surveys that are appropriate for evaluating the likely general effects of ORE development in any particular area. Tier 2 surveys are more detailed and correspond with archaeological surveys required by the Federal agency, Bureau of Ocean Energy Management (BOEM), prior to ORE development. The report also makes recommendations to improve the current BOEM guidelines and standards for archaeological surveys.

**Cultural and Historical Resources Task 2 – Cultural Landscape Approach (CLA).**

Task 2 provides the rationale and conceptual framework for CLA. Pioneered and partially implemented in the Ocean SAMP, the report uses RI waters as a case study. However, it is important to note CLA offers a multidisciplinary and multicultural approach to cultural heritage that operates along the full spectrum of geographic scales, from local to global.

The traditional way of assessing the impacts of coastal and offshore projects on cultural heritage resources involves focusing on the location, significance, and vulnerability of individual—physical—archaeological sites. Typically in the United States, this research has involved developing lists of the best-known shipwrecks and their possible locations with a particular project area. Frequently these lists or inventories are conjectural, as historically mentioned wrecks may or may not have actually occurred in a named area, or the wreck may have been recovered through unknown salvage operations. Despite a narrow focus, historic shipwrecks, unlike many underwater cultural resources, have at least received some consideration in coastal development, offshore oil and gas, and electrical and communication projects. They represent, however, only one of many kinds of potentially significant cultural heritage resources that might be adversely affected by ORE development. The recommendations in the report respond to widespread calls for the better integration of human factors in marine resource management, and for research and management methods that are sensitive to and inclusive of tribal and indigenous people and working maritime communities (Douvre 2008; Pomeroy and Douvre 2008; Crowder and Norse 2008; St. Thomas and Hall-Arbor 2008; Elher 2008).

CLA bridges traditional historic preservation-based approaches to maritime heritage resource management and the broader consideration and integration of human factors in the environment called for by EBM, Coastal and Marine Spatial Planning, and the National Ocean Policy. CLA addresses contemporary management challenges related to cultural heritage resources by providing an open-ended and rigorous framework that integrates data and perspectives from the physical and social sciences, humanities, and traditional/place-based knowledge systems. CLA is a holistic process that recognizes that places and cultural heritage resources can have different or multiple meanings and levels of significance based on how people from different cultures, times, or backgrounds have interacted with the landscape. Cultural heritage resources, whether in the form of archaeological sites or living cultural practices, are records of these interactions over time, and reveal how people have used and shaped marine environments, and how these environments have shaped human culture and history. Understanding these interactions may offer our best hope for sustainably and equitably using, maintaining, and where required restoring coastal and marine ecosystems (Crowder and Norse 2008; Douvre 2008). Adopting this

pluralistic approach increases the likelihood that cultural heritage resources will be found, recognized, and appropriately respected as decisions are made about issues, such as the siting and potential effects of ORE.

Task 2 of the report also aims to improve the performance of NEPA and Section 106 reviews under the National Historic Preservation Act (NHPA) and to bring them into better alignment with the National Ocean Policy and its nine priority areas. The report recommends the adoption of new definitions and categories for cultural heritage resources developed under the auspices of the National Marine Protected Area Federal Advisory Committee in 2010. The report also strongly recommends the need for early and meaningful consultation with tribal and indigenous groups, as well as members of working maritime communities in developing landscape contexts and preservation priorities.

**Cultural and Historical Resources Task 3 – Archaeological Sensitivity Analysis (ASA).**

Task 3 advocates for the use of ASA after comparing and evaluating three models (ASA, Predictive Modeling, and Paleo-Archaeological Landscape Reconstruction) in terms of potential effectiveness, practicality, and as a tool for ORE baseline studies for submerged cultural resources. ASA is a technique used by archaeologists and historians to designate certain areas as more archaeologically sensitive than others. In that sense it is closely allied to predictive modeling. Those designations are based on historic, archaeological, GIS, geophysical, and site-specific studies as interpreted by an experienced professional archaeologist and/or historian. All the data is geo-spatial in nature but not necessarily quantitative. ASA is not usually built upon a statistical model. More frequently, ASA is based on exploratory data analysis and is dependent on the capacity of the field professionals to see patterns, make judgments and divide an area into zones of archaeological sensitivity. Those zones tend to be 3 or 5 in number and range from *Highest Sensitivity* (areas that contain known cultural resources that are on, or have been determined eligible for inclusion in, the National Register of Historic Places) to *Lowest Sensitivity* (areas that have experienced low levels of documented human activity or that have experienced extensive disturbance. They contain no known historic or archaeological sites, a finding that has been confirmed through geophysical survey and archaeological inspection). Certainly ASA can and has been applied to submerged environments, for example by Mather and Watts in the James River and Charleston Harbor (Watts and Mather 1996; Watts and Mather

1997). In general, ASA has been used for assessments of historic rather than prehistoric site patterns and sensitivity (Mather and Watts 1998; Mather and Watts 2002).

Since ASA has the potential to identify areas with greater likelihood for containing archaeological resources, it can help developers and managers with assessment of time, costs and threats to cultural resources. A GIS-based ASA could also dovetail well with the results of paleo-archaeological landscape reconstruction.

Using the Ocean SAMP as a case study, the report attempts to refine and test ASA as a tool to predict historic sites (particularly shipwreck locations) and better explain wreck distributions using readily available data and linear regression. While the patterns of shipwreck loss revealed by the analysis of Rhode Island data may not be applicable in every location, the methodology for revealing those patterns is likely to be broadly pertinent.

**Public Feedback Received During the Public Comment Period**

Name	Affiliation	Comment	Response
Caroline Karp	Brown University	Need research to identify cumulative impacts from all offshore development	This is an important concept that had been indirectly conveyed through our emphasis on monitoring studies and taking an ecosystem perspective of the Ocean SAMP area. This concept has been more clearly expressed within the SRA in the "Introduction" and "Research Focused on Impact Assessment" sections.
Caroline Karp	Brown University	Emphasize how research can be used to identify conservation areas	This is an important point that has been made clearer throughout the SRA.
Caroline Karp	Brown University	Legal analysis must be conducted re: fixed private infrastructure on public trust lands	While this is a valid concern, it is beyond the scope of the SRA.
Kevin Ruddick	Nature Conservancy	Newport Naval Station and Nature Conservancy collaboration	The USN and TNC would be valuable collaborators for the research proposed in the SRA for the Ocean SAMP area and regionally. Section 6 "Regional Research Initiatives" now identifies USN and TNC as potential collaborators.
Kevin Ruddick	Nature Conservancy	identify other areas for [future?] conservation	The SRA is not a tool through which to identify areas in need of conservation. However, the SRA is a tool to propose research that will lead to the identification of such areas. The need to incorporate a conservation focus is important, and

we have made that clearer throughout the SRA.

Caroline Karp    Brown University

Research Agenda for inland, bay, etc.

While this is a valid concern, it is beyond the scope of this SRA.

Edward Rooney    Verizon

In the Ocean engineering section --stress the importance of these studies for CRMC and the nation to understand design and safety issues so better standards can be developed (e.g. CVA process, etc.)

The SRA mentioned the importance of engineering studies to better understand design and structural issues related to ORE development in order to improve standards. Based on this comment, these concepts were expanded upon in the ocean engineering section of the SRA and the important issue of safety was incorporated.

Edward Rooney    Verizon

must find out about zoning for industry, including lessons learned from land-based zoning

While this is a valid concern, it is beyond the scope of this SRA.

Richard Horwitz    (see pdf - submitted comments through Ocean SAMP comment form online)

I generally admire the physical science side of the agenda, but I also find the social science side extraordinarily narrow (basically exclusively economics with next to nothing from any of the other social sciences or their target media. E.g., "cultural" -- at least to anthropologists, sociologists, political scientists, and all but the most devout modelers -- means much

Understanding the social science aspects of the Ocean SAMP area has been a major focus of the Ocean SAMP and SRA processes. These efforts have expanded beyond only those related to economics and physical structures of the past. As an example, the recreation and tourism and the fisheries sections both deal with people's current perceptions. We feel like we presented a wide variety of methods to understand and assess

more than physical structures and the past. cultural realities and social conditions.  
In fact, the main predictors of conflict and conditions for compromise has had much more to do with people's current perceptions, even if they are not readily capture by naively objectivist quantitative methods. I'd urge consideration of attention to discovery of cultural realities and social conditions, much more broadly conceived and more flexibly assessed.

August 17, 2012

Grover Fugate

Executive Director

Rhode Island Coastal Resources Management Council

Via e-mail

Re: Comments for the RI Ocean SAMP Research Agenda

Dear Director Fugate,

The Nature Conservancy appreciates this opportunity to provide comments on the proposed research agenda for the Ocean Special Area Management Plan. In general, we support the draft agenda and feel the proposed research is essential to the success of the OSAMP.

It is important that the state continues to monitor living resources throughout the OSAMP area and integrates timely and up-to-date information on fish, marine mammals, avifauna, sea turtles, shellfish and crustaceans.

The plan highlights renewable energy and climate change impacts as important areas for study. We recommend a comprehensive scope that includes potential cumulative impacts of full build-out scenarios.

The entirety of human uses, now and into the foreseeable future need to be considered. The full range of activities in the OSAMP and across the region needs to be evaluated in the context of the cumulative impacts to the marine and coastal ecosystem.

The impacts of current human activities, including hydraulic dredging, scallop dredging and trawling need to be evaluated. It may be useful to identify which bottom types are most sensitive to specific

impacts. An approach similar to that used by the Nation Center for Ecological Analysis and Synthesis (NCEAS) could be used, considering the degree of impact as well as the sensitivity and resilience of sea floor habitats.

A synthesis and analysis of the existing and collected data with the purpose of identifying places and processes of high habitat value or sensitivity would be valuable. These analyses could help to identify areas for conservation; these could be ADPs (Areas Designated for Protection), APCs (Areas of Particular Concern) or a new class specific to the needs and sensitivities of the identified resource.

Provisions to measure public benefits, ecosystem services, economic benefits, and ecological benefits of protection and marine conservation should be included.

An explicit goal relating to the conservation of threatened and endangered species (Atlantic Sturgeon, common Loon, etc.) should be added.

Specific attention should be given to the health and viability of spawning habitat for various species, particularly areas with large concentrations of egg deposits.

Coastal habitats are an important part of the ecology of the Ocean SAMP. Consideration should be given to the inclusion of these places in the databases and analyses so that the system can better be evaluated in a holistic manner.

The classification and organization of data should coordinate with NOAA's Coastal and Marine Ecological Classification Standard (CMECS).

Study and consideration of the potential 'wake effect' downwind of turbines is needed. Water vapor 'contrails' could have negative impacts, particularly to Block Island and its airport.

Thank you for your consideration of these comments. If you have any questions, you may contact me directly at (401) 331-7110 x 12, or via e-mail at [kruddock@tnc.org](mailto:kruddock@tnc.org).

Sincerely,

A handwritten signature in blue ink, appearing to read "Kevin Ruddock". The signature is fluid and cursive, with the first name "Kevin" written in a larger, more prominent script than the last name "Ruddock".

Kevin Ruddock

GIS Analyst

# OSAMP Research Agenda Comment Form

#1

Name \* Richard Horwitz

Email \* rhorwitz@cox.net

Affiliation, Organization, or Town \* URI Coastal Institute and Independent Consultant

Please submit the comment template here.

In your opinion, which of the research initiatives listed in the OSAMP Research Agenda are the most important to pursue in the next 3-5 years? 3.0 . . . much more broadly conceived

When considering a 10-20 year timeline, does your answer change? If so, please explain. No.

### General comments on the Research Agenda:

I generally admire the physical science side of the agenda, but I also find the social science side extraordinarily narrow (basically exclusively economics with next to nothing from any of the other social sciences or their target media. E.g., "cultural" -- at least to anthropologists, sociologists, political scientists, and all but the most devout modelers -- means much more than physical structures and the past. In fact, the main predictors of conflict and conditions for compromise has had much more to do with people's current perceptions, even if they are not readily capture by naively objectivist quantitative methods. I'd urge consideration of attention to discovery of cultural realities and social conditions, much more broadly conceived and more flexibly assessed.

Created  
25 Jun 2012  
10:45:04 AM

68.9.117.69  
IP Address

PUBLIC